

GROWTH OF TORSION GROUPS OF ELLIPTIC CURVES UPON BASE CHANGE

ENRIQUE GONZÁLEZ–JIMÉNEZ AND FILIP NAJMAN

ABSTRACT. We study how the torsion of elliptic curves over number fields grows upon base change, and in particular prove various necessary conditions for torsion growth. For a number field F , we show that for a large set of number fields L , whose Galois group of their normal closure over F has certain properties, it will hold that $E(L)_{\text{tors}} = E(F)_{\text{tors}}$ for all elliptic curves E defined over F .

Our methods turn out to be particularly useful in studying the possible torsion groups $E(K)_{\text{tors}}$, where K is a number field and E is a base change of an elliptic curve defined over \mathbb{Q} . Suppose that E is a base change of an elliptic curve over \mathbb{Q} for the remainder of the abstract. We prove that $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$ for all elliptic curves E defined over \mathbb{Q} and all number fields K of degree d , where d is not divisible by a prime ≤ 7 . Using this fact, we determine all the possible torsion groups $E(K)_{\text{tors}}$ over number fields K of prime degree $p \geq 7$. We determine all the possible degrees of $[\mathbb{Q}(P) : \mathbb{Q}]$, where P is a point of prime order p for all p such that $p \not\equiv 8 \pmod{9}$ or $\left(\frac{-D}{p}\right) = 1$ for any $D \in \{1, 2, 7, 11, 19, 43, 67, 163\}$; this is true for a set of density $\frac{1535}{1536}$ of all primes and in particular for all $p < 3167$. Using this result, we determine all the possible prime orders of a point $P \in E(K)_{\text{tors}}$, where $[K : \mathbb{Q}] = d$ for all $d \leq 3342296$. Finally, we determine all the possible groups $E(K)_{\text{tors}}$, where K is a quartic number field and E is an elliptic curve defined over \mathbb{Q} and show that no quartic sporadic point on a modular curve $X_1(m, n)$ comes from an elliptic curve defined over \mathbb{Q} .

1. INTRODUCTION

Let E/F be an elliptic curve defined over a number field F . The Mordell-Weil theorem states that the set $E(F)$ of F -rational points is a finitely generated abelian group. Denote by $E(F)_{\text{tors}}$, the torsion subgroup of $E(F)$. It is well known that the possible torsion groups are of the shape $\mathcal{C}_m \times \mathcal{C}_n$ for two positive integers n, m , where m divides n and where \mathcal{C}_k is a cyclic group of order k .

The purpose of this paper is to shed light on how the torsion subgroup of an elliptic curve defined over a number field can grow when considered over an extension of the field of definition. We will mostly be interested in the case where L is also a number field, although some of our results apply to infinite extensions of F ; for example in Corollary 7.6 we prove that the torsion of elliptic curves over \mathbb{Q} does not grow in \mathbb{Z}_p -extensions of \mathbb{Q} for $p \geq 11$. Our results give necessary conditions

Received by the editor February 1, 2019, and, in revised form, May 22, 2019, and June 6, 2019.
2010 *Mathematics Subject Classification*. Primary 11G05.

Key words and phrases. Elliptic curves, torsion over number fields.

The first author was partially supported by the grant MTM2015–68524–P.

The second author gratefully acknowledges support from the QuantiXLie Center of Excellence and by the Croatian Science Foundation under the project no. IP-2018-01-1313.

on the degrees and Galois groups of extensions of number fields L/F depending on the growth from $E(F)_{\text{tors}}$ to $E(L)_{\text{tors}}$. The most general results of this type are proved in Section 4, where L/F is an arbitrary extension of number fields.

Naturally, we can obtain much more precise results if we specialize to the case of $F = \mathbb{Q}$. We will do this in Section 5. Suppose throughout the introduction that E is always defined over \mathbb{Q} . For $P \in E(\overline{\mathbb{Q}})$ define $\mathbb{Q}(P)$ to be the number field obtained by adjoining the coordinates of P to \mathbb{Q} . We study the problem of determining, for a given prime p , all the possible values of the degree of $[\mathbb{Q}(P) : \mathbb{Q}]$ for all points $P \in E(\overline{\mathbb{Q}})$ of order p on all elliptic curves E/\mathbb{Q} . This is done by studying the lengths of orbits of $E[p]$ under the action of the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. Here we rely heavily on the knowledge of the possible images $G_E(p)$ of mod p Galois representations attached to elliptic curves over \mathbb{Q} . While all the possibilities for $G_E(p)$ are not yet known in general, the state of the knowledge is good enough for our results, stated in Theorem 5.6, to be unconditional for all p such that $p \not\equiv 8 \pmod{9}$ or $\left(\frac{-D}{p}\right) = 1$ for any $D \in \{1, 2, 7, 11, 19, 43, 67, 163\}$. The set of primes satisfying these conditions has natural density $\frac{1535}{1536}$ and the smallest prime that does not satisfy these conditions is $p = 3167$. Should Serre’s uniformity Conjecture 3.3 or its weaker version, Conjecture 3.5, be proved our results will hold true for all primes p .

An important problem in the theory of elliptic curves is to characterize up to isomorphism the possible torsion groups of elliptic curves over a given number field, or over number fields of given degree. Before we state known results about this problem, we introduce some notation. For a given positive integer d ,

- Let $\Phi(d)$ be the set of possible isomorphism classes of groups $E(K)_{\text{tors}}$, where K runs through all number fields K of degree d and E runs through all elliptic curves over K .
- Let $\Phi_{\mathbb{Q}}(d) \subseteq \Phi(d)$ be the set of possible isomorphism classes of groups $E(K)_{\text{tors}}$, where K runs through all number fields K of degree d and E runs through all elliptic curves defined over \mathbb{Q} .
- Let $\Phi^{\infty}(d)$ be the subset of isomorphism classes of groups $\Phi(d)$ that occur infinitely often. More precisely, a torsion group G belongs to $\Phi^{\infty}(d)$ if there are infinitely many elliptic curves E , non-isomorphic over $\overline{\mathbb{Q}}$, such that $E(K)_{\text{tors}} \simeq G$.
- Let $R_{\mathbb{Q}}(d)$ be the set of all primes p such that there exists a number field K of degree d , an elliptic curve E/\mathbb{Q} such that there exists a point of order p on $E(K)$.

Mazur [29] determined $\Phi(1)$, while Kamienny, Kenku, and Momose [20, 25] determined $\Phi(2)$. For $d \geq 3$, the set $\Phi(d)$ is, at the moment of the writing of the paper, not known. Merel [31] proved that $\Phi(d)$ is finite for all positive integers d .

It is trivial to see that $\Phi(1) = \Phi^{\infty}(1)$ and $\Phi(2) = \Phi^{\infty}(2)$; this follows immediately from the fact that all the modular curves that parameterize the groups from $\Phi(1)$ are of genus 0 and that the modular curves that parameterize the groups from $\Phi(2)$ are of genus ≤ 2 . The set $\Phi^{\infty}(d)$ has been determined for $d = 3$ by Jeon, Kim, and Schweizer [17]; for $d = 4$ by Jeon, Kim, and Park [18]; and for $d = 5$ and 6, recently, by Derickx and Sutherland [10].

In [32] the second author determined $\Phi_{\mathbb{Q}}(2)$ and $\Phi_{\mathbb{Q}}(3)$. In this paper we determine $\Phi_{\mathbb{Q}}(p)$ for all primes $p \geq 7$ (see Corollary 7.3). Together with the paper [11]

of the first author, which determines $\Phi_{\mathbb{Q}}(5)$, this completes the determination of all possible torsion groups over prime degree number fields of elliptic curves defined over \mathbb{Q} .

This paper is, as far as we are aware, the first to study the set $R_{\mathbb{Q}}(d)$. Lozano-Robledo [28] determined the sets $S_{\mathbb{Q}}(d) = \bigcup_{k \leq d} R_{\mathbb{Q}}(k)$ for $d \leq 42$. Note that the knowledge of $R_{\mathbb{Q}}(d)$ for all $d \leq B$ obviously gives $S_{\mathbb{Q}}(d)$ for all $d \leq B$, for any bound B , but not vice versa. Hence $R_{\mathbb{Q}}(d)$ can be considered to be a somewhat "finer" invariant. In Section 6 we determine $R_{\mathbb{Q}}(d)$ for all $d \leq 3342296$. This is done by applying the results of Section 5.

In Section 7 we use the results of Section 5 to show, in Theorem 7.2, that $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$ for all number fields K of degree d , where d is an integer whose smallest prime divisor is ≥ 11 . In particular this applies for all number fields of prime degree p for $p \geq 11$.

In [32] it was shown that $\Phi_{\mathbb{Q}}(3) \not\subseteq \Phi^{\infty}(3)$, which implies $\Phi(3) \neq \Phi^{\infty}(3)$. Derickx and van Hoeij [9] proved that $\Phi(n) \neq \Phi^{\infty}(n)$ for $5 \leq n \leq 17$. Thus, it is natural to ask whether $\Phi(4) = \Phi^{\infty}(4)$. It was proved that $\Phi(3) \neq \Phi^{\infty}(3)$ by showing $\Phi_{\mathbb{Q}}(3) \not\subseteq \Phi^{\infty}(3)$, thus it is natural to try to determine $\Phi_{\mathbb{Q}}(4)$, apart from it being interesting in its own right, for this purpose. This problem, of trying to determine $\Phi_{\mathbb{Q}}(4)$, originally motivated this work. We solve this problem completely in Section 8 and obtain that $\Phi_{\mathbb{Q}}(4) \subset \Phi^{\infty}(4)$.

Some of the proofs in the paper rely on extensive computations in Magma [5]. This primarily applies to computations in fixed finite groups. Where it is obvious that we are doing a computation in a fixed finite group, we will do it often without mention. All of the programs and calculation used for the proofs can be found at the research website of the first author [14].

2. NOTATION AND AUXILIARY RESULTS

In this section we fix notation throughout the paper. If E is an elliptic curve, then j_E will denote its j -invariant. We will say that E has CM or is a CM elliptic curve if it has complex multiplication. Recall that an elliptic curve E has complex multiplication if its endomorphism ring is isomorphic to an order of a quadratic imaginary field. If it does not have complex multiplication, we say that the curve is non-CM.

There exists an F -rational cyclic isogeny $\phi : E \rightarrow E'$ of degree n if and only if $\ker \phi$ is a $\text{Gal}(\overline{F}/F)$ -invariant cyclic group of order n ; in this case we say that E/F has an n -isogeny.

In this section we will list, for the sake of the reader, some known results that we will be using throughout the paper.

The possible degrees of n -isogenies of elliptic curves over \mathbb{Q} are known by the following theorem.

Theorem 2.1 (Mazur [30] and Kenku [21–24]). *Let E/\mathbb{Q} be an elliptic curve with an n -isogeny over \mathbb{Q} . Then*

$$(1) \quad n \in \{1, \dots, 19, 21, 25, 27, 37, 43, 67, 163\}.$$

There are infinitely many elliptic curves (up to $\overline{\mathbb{Q}}$ -isomorphism) with an n -isogeny over \mathbb{Q} for $n \in \{1, \dots, 10, 12, 13, 16, 18, 25\}$ and only finitely many for all the other n listed in (1).

Furthermore, if $n \in \{14, 19, 27, 43, 67, 163\}$ and E has an n -isogeny over \mathbb{Q} , then E has CM.

Mazur [29] determined $\Phi(1)$:

$$\Phi(1) = \{\mathcal{C}_n : n = 1, \dots, 10, 12\} \cup \{\mathcal{C}_2 \times \mathcal{C}_{2n} | n = 1, \dots, 4\}.$$

Kamienny [20] and Kenku and Momose [25] determined $\Phi(2)$:

$$\begin{aligned} \Phi(2) = \{\mathcal{C}_n : n = 1, \dots, 16, 18\} \cup \{\mathcal{C}_2 \times \mathcal{C}_{2n} | n = 1, \dots, 6\} \\ \cup \{\mathcal{C}_3 \times \mathcal{C}_{3n} | n = 1, 2\} \cup \{\mathcal{C}_4 \times \mathcal{C}_4\}. \end{aligned}$$

The second author [32] determined $\Phi_{\mathbb{Q}}(2)$:

$$\Phi_{\mathbb{Q}}(2) = \{\mathcal{C}_n : n = 1, \dots, 10, 12, 15, 16\} \cup \{\mathcal{C}_2 \times \mathcal{C}_{2n} : n = 1, \dots, 6\}.$$

3. IMAGES OF MOD n GALOIS REPRESENTATIONS

When studying the growth of torsion groups upon extensions, one is naturally led to the study of mod n Galois representations attached to elliptic curves. Let E be an elliptic curve over a number field K and let n be a positive integer. We denote by $E[n]$ the n -torsion subgroup of $E(\overline{K})$, where \overline{K} is a fixed algebraic closure of K . That is, $E[n] = \{P \in E(\overline{K}) \mid [n]P = \mathcal{O}\}$, where \mathcal{O} denotes the identity of the group $E(\overline{K})$. The field $K(E[n])$ is the number field obtained by adjoining all the x - and y -coordinates of the points of $E[n]$, or, equivalently, the smallest field over which all of $E[n]$ is defined. The absolute Galois group $\text{Gal}(\overline{K}/K)$ acts on $E[n]$ by its action on the coordinates of the points, inducing a *mod n Galois representation attached to E*

$$\rho_{E,n} : \text{Gal}(\overline{K}/K) \longrightarrow \text{Aut}(E[n]).$$

Notice that since $E[n]$ is a free $\mathbb{Z}/n\mathbb{Z}$ -module of rank 2, fixing a basis $\{P, Q\}$ of $E[n]$, we identify $\text{Aut}(E[n])$ with $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$. Then we rewrite the above Galois representation as

$$\rho_{E,n} : \text{Gal}(\overline{K}/K) \longrightarrow \text{GL}_2(\mathbb{Z}/n\mathbb{Z}).$$

Therefore we can view $\rho_{E,n}(\text{Gal}(\overline{K}/K))$ as a subgroup of $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$, determined uniquely up to conjugacy in $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$, and denoted by $G_E(n)$ from now on; it will always be clear what the field K is. In certain instances we will say that $G_E(n) = G$ for some group G ; this will always mean that we have fixed a basis so that we remove ambiguity of working up to conjugacy.

Since $K(E[n]) = \{x, y \mid (x, y) \in E[n]\}$ is Galois over K and since $\ker \rho_{E,n} = \text{Gal}(\overline{K}/K(E[n]))$, we deduce that $G_E(n) \simeq \text{Gal}(K(E[n])/K)$. Let $R = (x(R), y(R)) \in E[n]$ and $K(R) = K(x(R), y(R)) \subseteq K(E[n])$. Then by Galois theory there exists a subgroup \mathcal{H}_R of $\text{Gal}(K(E[n])/K)$ such that $K(R) = K(E[n])^{\mathcal{H}_R}$. In particular, if we denote by H_R the image of \mathcal{H}_R in $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$, we have:

- $[K(R) : K] = [G_E(n) : H_R]$,
- $\text{Gal}(\widehat{K(R)}/K) \simeq G_E(n)/\mathcal{N}_{G_E(n)}(H_R)$, where $\widehat{K(R)}$ denotes the Galois closure of $K(R)$ over K , and $\mathcal{N}_{G_E(n)}(H_R)$ denotes the normal core of H_R in $G_E(n)$.

We will use the following obvious lemma, sometimes without mention.

Lemma 3.1. *Let E be an elliptic curve over a number field K , let n be a positive integer, and $R \in E[n]$. Then $[K(R) : K]$ divides $|G_E(n)|$. In particular $[K(R) : K]$ divides $|\text{GL}_2(\mathbb{Z}/n\mathbb{Z})|$.*

In practice, given the conjugacy class of $G_E(n)$ in $\text{GL}_2(\mathbb{Z}/n\mathbb{Z})$, we can deduce the relevant arithmetic properties of the fields of definition of the n -torsion points: since $E[n]$ is a free $\mathbb{Z}/n\mathbb{Z}$ -module of rank 2, we can identify the n -torsion points with $(a, b) \in (\mathbb{Z}/n\mathbb{Z})^2$ (i.e., if $R \in E[n]$ and $\{P, Q\}$ is a $\mathbb{Z}/n\mathbb{Z}$ -basis of $E[n]$, then there exist $a, b \in \mathbb{Z}/n\mathbb{Z}$ such that $R = aP + bQ$). Therefore H_R is the stabilizer of (a, b) by the action of $G_E(n)$ on $(\mathbb{Z}/n\mathbb{Z})^2$.

Let p be an odd prime and $\epsilon = -1$ if $p \equiv 3 \pmod{4}$ and otherwise let $\epsilon \geq 2$ be the smallest integer such that $\binom{\epsilon}{p} = -1$.

Define the following matrices in $\text{GL}_2(\mathbb{F}_p)$ for some $a, b \in \mathbb{F}_p$:

$$D(a, b) = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \quad M_\epsilon(a, b) = \begin{pmatrix} a & b\epsilon \\ b & a \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \text{and} \quad J = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

We define the following subgroups of $\text{GL}_2(\mathbb{F}_p)$:

$$\begin{aligned} C_s(p) &= \{D(a, b) : a, b \in \mathbb{F}_p^\times\}, \\ C_s^+(p) &= \{D(a, b), T \cdot D(a, b) : a, b \in \mathbb{F}_p^\times\}, \\ C_{ns}(p) &= \{M_\epsilon(a, b) : (a, b) \in \mathbb{F}_p^2, (a, b) \neq (0, 0)\}, \\ C_{ns}^+(p) &= \{M_\epsilon(a, b), J \cdot M_\epsilon(a, b) : (a, b) \in \mathbb{F}_p^2, (a, b) \neq (0, 0)\}. \end{aligned}$$

The group $C_s(p)$ is called the *split Cartan subgroup* and $C_{ns}(p)$ is called the *non-split Cartan subgroup*. Note that $C_s^+(p)$ (resp., $C_{ns}^+(p)$) is the normalizer of $C_s(p)$ (resp., $C_{ns}(p)$) in $\text{GL}_2(\mathbb{F}_p)$. The orders of the groups $C_s(p)$, $C_s^+(p)$, $C_{ns}(p)$, and $C_{ns}^+(p)$ are $(p - 1)^2$, $2(p - 1)^2$, $p^2 - 1$, and $2(p^2 - 1)$, respectively.

Notation. Conjugacy classes of subgroups of $\text{GL}_2(\mathbb{F}_p)$, for fixed small values of p , will be identified in this paper by labels introduced by Sutherland [36, §6.4] and used in the LMFDB database [27]. We should note that Zywina [37] uses different notation for such conjugacy classes of subgroups of $\text{GL}_2(\mathbb{F}_p)$. To aid the reader, in Tables 1 and 2 we list both Sutherland’s and Zywina’s names for each of the groups.

We turn our attention to what is known on the possible images of mod p Galois representations attached to elliptic curves defined over \mathbb{Q} , where p is a prime. Determining all the possibilities of the images $G_E(p) \subseteq \text{GL}_2(\mathbb{F}_p)$ for elliptic curves defined over \mathbb{Q} is an important (and still open) problem in the theory of elliptic curves. The following theorem, due to Mazur, Serre, Bilu, Parent, Rebolledo, Zywina, Balakrishnan, Dogra, Müller, Tuitman, and Vonk (see [2–4, 30, 34, 37] for more details) gives the current state of the knowledge for non-CM elliptic curves.

Theorem 3.2. *Let E/\mathbb{Q} be a non-CM elliptic curve and let p be a prime. Then one of the following possibilities occurs:*

- (i) $G_E(p) = \text{GL}_2(\mathbb{F}_p)$.
- (ii) $p \in \{2, 3, 5, 7, 11, 13, 17, 37\}$, and $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to one of the groups in Tables 1 and 2.
- (iii) $p \geq 17$:
 - (a) $p \equiv 1 \pmod{3}$, and $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_{ns}^+(p)$.
 - (b) $p \equiv 2 \pmod{3}$, and $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_{ns}^+(p)$ or to the subgroup

$$G_0(p) := \{M_\epsilon(a, b)^3, J \cdot M_\epsilon(a, b)^3 : (a, b) \in \mathbb{F}_p^2, (a, b) \neq (0, 0)\} \subseteq C_{ns}^+(p).$$

The complete and unconditional description of possible images $G_E(p)$ for $p \leq 11$ appearing in Theorem 3.2, which are compiled in Tables 1 and 2 in the appendix, is due to Zywna [37] (see also Sutherland’s paper [36]).

Before going any further we mention an important open conjecture, motivated by a question by Serre [35] (see also [37, Conjecture 1.12.]), which conjecturally characterizes the possible images of Galois representations attached to non-CM elliptic curves.

Conjecture 3.3. *If E is a non-CM elliptic curve defined over \mathbb{Q} , $p \geq 17$ a prime, and (p, j_E) not in the set*

$$(2) \quad \{(17, -17 \cdot 373^3/2^{17}), (17, -17^2 \cdot 101^3/2), (37, -7 \cdot 11^3), (37, -7 \cdot 137^3 \cdot 2083^3)\},$$

then $G_E(p) = \text{GL}_2(\mathbb{F}_p)$.

Remark 3.4. If Conjecture 3.3 is true, the case (iii) does not occur. All of the other cases are known to occur.

Some of our results will be best possible under the following conjecture, which is a weaker version of Conjecture 3.3.

Conjecture 3.5. *If E is a non-CM elliptic curve defined over \mathbb{Q} , $p \geq 17$ a prime, and (p, j_E) not in the set*

$$(3) \quad \{(17, -17 \cdot 373^3/2^{17}), (17, -17^2 \cdot 101^3/2), (37, -7 \cdot 11^3), (37, -7 \cdot 137^3 \cdot 2083^3)\},$$

then $G_E(p) = \text{GL}_2(\mathbb{F}_p)$ or $C_{ns}^+(p)$.

The difference between Conjectures 3.3 and 3.5 is that in the latter we allow that $G_E(p) = C_{ns}^+(p)$. For our purposes, there is no difference between $C_{ns}^+(p)$ and $\text{GL}_2(\mathbb{F}_p)$, as both act transitively on the non-zero elements of \mathbb{F}_p^2 .

If E/\mathbb{Q} is a CM elliptic curve and p a prime, the theory of complex multiplication gives us a lot of information about $G_E(p)$. In the CM case, the possibilities for $G_E(p)$ are completely understood. We list the possibilities for $G_E(p)$ (in the form stated in [37, §1.9]).

Theorem 3.6. *Let E/\mathbb{Q} be a CM elliptic curve and let p be a prime. The ring of endomorphisms of $E_{\overline{\mathbb{Q}}}$ is an order of conductor f in the ring of integers of an imaginary quadratic field of discriminant $-D$.*

- (i) *If $p = 2$, then $G_E(2) = \text{GL}_2(\mathbb{F}_2)$ or is conjugate in $\text{GL}_2(\mathbb{F}_2)$ to $2B$ or $2Cs$.*
- (ii) *If $p > 2$ and $(D, f) \neq (3, 1)$:*
 - (a) *If $(\frac{-D}{p}) = 1$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_s^+(p)$.*
 - (b) *If $(\frac{-D}{p}) = -1$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_{ns}^+(p)$.*
 - (c) *Suppose that p divides D and hence $D = p$. Then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to one of the following subgroups:*

$$G_{00}(p) := \left\{ \begin{pmatrix} a & b \\ 0 & \pm a \end{pmatrix} : a \in \mathbb{F}_p^\times, b \in \mathbb{F}_p \right\},$$

$$G_{10}(p) := \left\{ \begin{pmatrix} a & b \\ 0 & \pm a \end{pmatrix} : a \in (\mathbb{F}_p^\times)^2, b \in \mathbb{F}_p \right\},$$

or

$$G_{01}(p) := \left\{ \begin{pmatrix} \pm a & b \\ 0 & a \end{pmatrix} : a \in (\mathbb{F}_p^\times)^2, b \in \mathbb{F}_p \right\}.$$

- (iii) If $p > 2$ and $(D, f) = (3, 1)$:
 - (a) If $p \equiv 1 \pmod{9}$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_s^+(p)$.
 - (b) If $p \equiv 8 \pmod{9}$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_{ns}^+(p)$.
 - (c) If $p \equiv 4$ or $7 \pmod{9}$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_s^+(p)$ or to the subgroup

$$G^3(p) := \{D(a, ab^3), T \cdot D(a, ab^3) : a, b \in \mathbb{F}_p^\times\} \subseteq C_s^+(p).$$
 - (d) If $p \equiv 2$ or $5 \pmod{9}$, then $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to $C_{ns}^+(p)$ or to the subgroup $G_0(p)$ of $C_{ns}^+(p)$.
 - (e) If $p = 3$, then $G_E(3)$ is conjugate in $\text{GL}_2(\mathbb{F}_3)$ to $3Cs.1.1$, $3Cs$, $3B.1.1$, $3B.1.2$ or $3B$.

All the cases occur.

4. GROWTH OF TORSION IN EXTENSIONS

In this section we prove general results, without any constraints on the base field over which the elliptic curve is defined, concerning the growth of the torsion of elliptic curves upon base change. We believe the results are primarily interesting in their own right, but they will also be useful in Section 8.

Throughout this section, unless otherwise specified, F is a number field, E/F an elliptic curve, and L/F is a finite extension with normal closure \widehat{L} .

Our first result gives restrictions on the growth of the p -torsion, in terms of the Galois group of the extension.

Theorem 4.1. *Let $G = \text{Gal}(\widehat{L}/F)$, and suppose that $H = \text{Gal}(\widehat{L}/L)$ is a non-normal maximal subgroup of G . Let p be a prime, $a = [F(\zeta_p) : F]$, and suppose G does not contain a cyclic quotient group of order a . Then $E(L)[p] = E(F)[p]$.*

Proof. Suppose that $P \in E(L)$, $P \notin E(F)$ is a point of order p . Then, as H is not normal in G there exists a $\sigma \in G$ such that $\sigma(L) \neq L$. It follows that $P^\sigma \in (E(L))^\sigma = E(\sigma(L))$. Note that as H is not contained in any proper subgroup of G , it follows that L has no proper subfield containing F , so $L \cap L^\sigma = F$. As P and P^σ are defined over different fields, it follows that they are independent, i.e., $\langle P, P^\sigma \rangle = E[p]$. Obviously, $P, P^\sigma \in E(\widehat{L})$, so $E(\widehat{L})[p] \simeq \mathcal{C}_p \times \mathcal{C}_p$. But this is impossible, since \widehat{L} does not contain ζ_p , because $F(\zeta_p)$ would be a cyclic extension of F of degree a , contradicting our assumptions. \square

By specializing the above theorem, we immediately obtain the following corollary.

Corollary 4.2. *Let E/\mathbb{Q} be an elliptic curve and let L be a number field with Galois closure \widehat{L} over \mathbb{Q} and with $G := \text{Gal}(\widehat{L}/\mathbb{Q})$. Suppose there are no intermediate fields $L \supseteq M \supseteq \mathbb{Q}$. Then:*

- (i) *If G is a simple group, and $p \geq 3$ is a prime, then $E(L)[p] = E(\mathbb{Q})[p]$.*
- (ii) *If $G \simeq A_n$ for $n \geq 4$, and $p \geq 3$ is a prime, then $E(L)[p] = E(\mathbb{Q})[p]$.*
- (iii) *If $G \simeq S_n$ for $n \geq 4$, and $p \geq 5$ is a prime, then $E(L)[p] = E(\mathbb{Q})[p]$.*

Proof. Statement (i) obviously follows from Theorem 4.1, and statement (ii) follows from the fact that A_n is simple for $n \geq 5$. It is easy to check that the conditions of Theorem 4.1 are satisfied for A_4 .

Statement (iii) follows for $n \geq 5$, from the fact that only a proper normal subgroup of S_n is A_n . For $n = 4$, it is again easy to check that there are no cyclic quotients of order ≥ 4 of S_4 . \square

We give another result that is useful for proving that over an extension with a certain Galois group there cannot be torsion growth of certain type.

Theorem 4.3. *Suppose L/F has no intermediate fields, and let $G = \text{Gal}(\widehat{L}/F)$. If G is not isomorphic to a quotient of $\text{Gal}(F(E[p])/F)$, then $E(L)[p] = E(F)[p]$.*

Proof. Suppose the opposite, that $E(F)[p] \subsetneq E(L)[p]$. It follows, since $L \supseteq F$ has no intermediate fields, that $L \subseteq F(E[p])$. Since $F(E[p])$ is a Galois extension of F , it follows that $\widehat{L} \subseteq F(E[p])$, since \widehat{L} is contained in any Galois extension of F which contains L . It follows that G is a quotient of $\text{Gal}(F(E[p])/F)$. \square

Remark 4.4. If $\text{Gal}(F(E[p])/F)$ is a priori unknown, one can still use the fact that $\text{Gal}(F(E[p])/F)$ is always a subgroup of $\text{GL}_2(\mathbb{F}_p)$. Then, in the setting as above, if G is not a quotient of any subgroup of $\text{GL}_2(\mathbb{F}_p)$, then $E(L)[p] = E(F)[p]$.

The previous two theorems give constraints on the growth of $E[p]$. One is naturally led to consider whether there can be any p^{n+1} torsion over an extension L of F if $E(F)[p] \neq \{\mathcal{O}\}$. The following result will be useful.

Theorem 4.5. *Let $G = \text{Gal}(\widehat{L}/F)$, let n be a positive integer, and let p be a prime not dividing $[L : F]$. Suppose that G is not isomorphic to a quotient of any subgroup of $\text{GL}_2(\mathbb{Z}/p^n\mathbb{Z})$ and that $\text{Gal}(\widehat{L}/L)$ is maximal in G . Let E/F be an elliptic curve such that it has an F -rational point of order p^n , but no F -rational points of order p^{n+1} . Then $E(L)$ has no points of order p^{n+1} .*

Proof. Suppose that there exists a point P of order p^{n+1} in $E(L)$. Since $\text{Gal}(\widehat{L}/L)$ is maximal in G , it follows that $L \supseteq F$ has no intermediate fields, so L is the field of definition of P , i.e., $L = F(P)$.

If $L \subseteq F(E[p^n])$ was true, then $\text{Gal}(F(E[p^n])/L)$ would be a quotient of $G_E(p^n) \simeq \text{Gal}(F(E[p^n])/F)$, which is in turn a subgroup of $\text{GL}_2(\mathbb{Z}/p^n\mathbb{Z})$, contradicting our assumptions. We conclude that $L \not\subseteq F(E[p^n])$.

Note that since $G_E(p^{n+1})$ is a subgroup of the inverse image of reduction mod p^n to $G_E(p^n)$, it follows that $[F(E[p^{n+1}]) : F(E[p^n])]$ divides p^4 . Since $P \in F(E[p^{n+1}])$ and $P \notin F(E[p^n])$, it follows that $F(E[p^n]) \subsetneq F(E[p^{n+1}])$. We conclude that $[F(E[p^{n+1}]) : F(E[p^n])] = p^k$, where $k = 1, 2, 3$, or 4 .

Now since both $L = F(P)$ and $F(E[p^n])$ are contained in $F(E[p^{n+1}])$, it follows that

$$F(E[p^n]) \subseteq LF(E[p^n]) \subseteq F(E[p^{n+1}]).$$

Since $L \not\subseteq F(E[p^n])$, we conclude that $[LF(E[p^n]) : F(E[p^n])]$ divides $[F(E[p^{n+1}]) : F(E[p^n])]$ and is hence equal to p^s where $s = 1, 2, 3$, or 4 . But since $[L : F]$ is coprime to $[F(E[p^n]) : F]$, we have

$$[LF(E[p^n]) : F(E[p^n])][F(E[p^n]) : F] = [LF(E[p^n]) : F] = [L : F][F(E[p^n]) : F],$$

so $[L : F] = [LF(E[p^n]) : F(E[p^n])]$, which obviously cannot be true since the left and right hand sides are coprime and > 1 . \square

Proposition 4.6. *Let n be a positive integer and let $P \in E(\overline{F})$ be a point of order p^{n+1} . Then $[F(P) : F(pP)]$ divides p^2 or $(p - 1)p$.*

Proof. Let $G = \text{Gal}(F(E[p^{n+1}])/F(pP))$, and $H = \text{Gal}(F(E[p^{n+1}])/F(P))$. By Galois theory, it follows that $[F(P) : F(pP)] = [G : H]$. To remove the ambiguity of having everything up to conjugacy in $\text{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z})$, let $\{P, Q\}$ be a basis of

$E[p^{n+1}]$ for some $Q \in E[p^{n+1}]$, let $G_E(p^{n+1})$ be induced by the action of $\text{Gal}(\overline{F}/F)$ on this basis $\{P, Q\}$, and consider G and H to be subgroups of $G_E(p^{n+1})$. Then in particular, G is a subgroup of

$$(4) \quad G' := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}) : a - 1 \equiv c \equiv 0 \pmod{p^n} \right\}.$$

Moreover, $|G|$ divides $p^2 \cdot p^n(p^n - p^{n-1}) = p^{2n+1}(p - 1)$. Now let

$$\Gamma = \left\{ \begin{pmatrix} 1 & x \\ 0 & y \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}) \right\};$$

it follows that H is isomorphic to $\Gamma \cap G$. Let B be the kernel of reduction of $\text{GL}_2(\mathbb{Z}/p^{n+1}\mathbb{Z}) \pmod{p^n}$. We have $|B| = p^4$. Now we have that the kernel of reduction mod p^n restricted to G (resp., H) is $B \cap G$ (resp., $B \cap H$). Since $B \cap H$ and $B \cap G$ are subgroups of B , it follows that their order is a power of p . We see that $H \pmod{p^n}$ is a subgroup of $G \pmod{p^n}$ of index d (where d divides $|\mathcal{G} \pmod{p^n}|$), so

$$|G \pmod{p^n}| = d \cdot |H \pmod{p^n}|, \quad \text{where } d|(p - 1)p^{2n-1}.$$

So

$$|G|/|B \cap G| = |G \pmod{p^n}| = d \cdot |H \pmod{p^n}| = d \cdot |H|/|B \cap H|,$$

and

$$|G|/|H| = d \cdot |B \cap G|/|B \cap H|.$$

The right hand side of the equations can be divisible only by powers of p and by divisors of $p - 1$. From the definitions of G and H , it follows that $[G : H] \leq p^2$, proving the proposition. \square

Remark 4.7. The results of Proposition 4.6 are best possible, in the sense that there exist both E/\mathbb{Q} and $P \in E(\overline{\mathbb{Q}})$ such that $[\mathbb{Q}(P) : \mathbb{Q}(pP)] = p^2$ and $[\mathbb{Q}(P) : \mathbb{Q}(pP)] = p(p - 1)$. The first case is “generic” and happens for any point of p -power order for any elliptic curve such that the p -adic representation attached to E is surjective.

The case when $[\mathbb{Q}(P) : \mathbb{Q}(pP)] = p(p - 1)$ occurs for example when $G_E(p^2)$ is the reduction of $\Gamma_0(p^2) \cup \Gamma_1(p) \pmod{p^2}$. Then $E(\mathbb{Q})$ contains a point Q of order p , and some of the solutions of $pP = Q$ are defined over a degree $p(p - 1)$ extension (and some over a degree p extension). One does not have to look far for an explicit example of such an elliptic curve—the elliptic curve 11a3 in the LMFDB [27] is such a curve for $p = 5$.

Proposition 4.8. *Let n be a positive integer, let $P \in E(\overline{F})$ be a point of order 2^{n+1} , and let $\widehat{F(P)}$ be the Galois closure of $F(P)$ over $F(2P)$. Then $[F(P) : F(2P)]$ divides 4 and $\text{Gal}(\widehat{F(P)}/F(2P))$ is either trivial or isomorphic to \mathcal{C}_2 , $\mathcal{C}_2 \times \mathcal{C}_2$, or D_4 .*

Proof. Let $G = \text{Gal}(F(E[2^{n+1}])/F(2P))$ and $H = \text{Gal}(F(E[2^{n+1}])/F(P))$. Then by Galois theory, $N = \text{Gal}(F(E[2^{n+1}])/\widehat{F(P)})$ is the normal core of H in G (i.e., the intersection of all conjugates of H by elements of G) and

$$\text{Gal}(\widehat{F(P)}/F(2P)) \simeq G/N.$$

To remove the ambiguity of having everything up to conjugacy in $\text{GL}_2(\mathbb{F}_p)$, let $\{P, Q\}$ be a basis of $E[2^{n+1}]$ for some $Q \in E[2^{n+1}]$, let $G_E(2^{n+1})$ be induced by

the action of $\text{Gal}(\overline{F}/F)$ on the basis $\{P, Q\}$, and consider G and H to be subgroups of $G_E(2^{n+1})$. We have that G is a subgroup of the group

$$(5) \quad G' = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/2^{n+1}\mathbb{Z}) : a - 1 \equiv c \equiv 0 \pmod{2^n} \right\}.$$

Define

$$\Gamma = \left\{ \begin{pmatrix} 1 & x \\ 0 & y \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/2^{n+1}\mathbb{Z}) \right\}.$$

Then H is isomorphic to $G \cap \Gamma$. We see that H is a subgroup of G whose index divides 4, so we obtain that $[F(P) : \widehat{F(2P)}]$ divides 4.

If $F(P) = F(2P)$, then obviously $\widehat{F(P)} = F(P) = F(2P)$, and the statement of the proposition is true.

If $[F(P) : F(2P)] = 2$, then obviously $F(P)$ is Galois over $F(2P)$ and

$$\text{Gal}(\widehat{F(P)}/F(2P)) \simeq \mathcal{C}_2.$$

If $[F(P) : F(2P)] = 4$, then it follows that

$$G/H = \left\{ H, \begin{pmatrix} 1 & 0 \\ 2^n & 1 \end{pmatrix} H, \begin{pmatrix} 1 + 2^n & 0 \\ 2^n & 1 \end{pmatrix} H, \begin{pmatrix} 1 & 0 \\ 2^n & 1 \end{pmatrix} H \right\}.$$

As N is the intersection of the elements of G/H , by direct calculation we obtain that

$$N = H \cap \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Z}/2^{n+1}\mathbb{Z}) : a - 1 \equiv c \equiv 0 \pmod{2^{n+1}}, b \equiv 0 \pmod{2} \right\}.$$

There are now 2 possibilities: either $N = H$ or $[H : N] = 2$. If $N = H$, then hence $F(P)$ is Galois over $F(2P)$, and we obtain that $G/N \simeq \mathcal{C}_2 \times \mathcal{C}_2$. On the other hand, if $[H : N] = 2$, then $G/N \simeq D_4$. □

5. DEGREE OF THE FIELD OF DEFINITION OF p -TORSION POINTS ON ELLIPTIC CURVES OVER \mathbb{Q}

In this section, for any prime p we determine all the possible degrees $[\mathbb{Q}(P) : \mathbb{Q}]$ for P a point of order p on an elliptic curve E/\mathbb{Q} .

Let E/\mathbb{Q} be an elliptic curve and let p be a prime. Then $G_E(p) \subseteq \text{GL}_2(\mathbb{F}_p)$ acts on $E[p] \simeq \mathbb{F}_p^2$. To remove any ambiguity, we choose that $G_E(p)$ acts on \mathbb{F}_p^2 from the left. The set \mathbb{F}_p^2 can be written as the disjoint union of the orbits of \mathbb{F}_p^2 by $G_E(p)$, and by the Orbit-Stabilizer Theorem we obtain

$$(6) \quad \#\mathbb{F}_p^2 = p^2 = \sum_{i=1}^k [\mathbb{Q}(R_i) : \mathbb{Q}] = \sum_{i=1}^k |G_E(p) \cdot (a_i, b_i)|, \quad \text{where } R_i = a_i P + b_i Q,$$

$(a_1, b_1), \dots, (a_k, b_k)$ are representatives of the orbits of \mathbb{F}_p^2 by $G_E(p)$, and $G_E(p) \cdot v$ denotes the orbit of $v \in \mathbb{F}_p^2$ by the action of $G_E(p)$.

In order to determine all the possible degrees of the fields of definition of the p -torsion points of E , by equation (6), we only need to know $G_E(p) \subseteq \text{GL}_2(\mathbb{F}_p)$ and the cardinality of the orbits of \mathbb{F}_p^2 by $G_E(p)$.

If we want to determine all the possible degrees of the extensions $\mathbb{Q}(P)$ over \mathbb{Q} for points $P \in E(\overline{\mathbb{Q}})$ of order p for any elliptic curve E/\mathbb{Q} we must, for all possible $G_E(p)$ and all $v \in \mathbb{F}_p^2, v \neq (0, 0)$, determine the index $[G_E(p) : G_E(p)_v]$ (where $G_E(p)_v$ is the stabilizer of v by the action of $G_E(p)$ on \mathbb{F}_p^2), or equivalently

$|G_E(p).v|$. Thanks to Theorems 3.2 and 3.6 we know a set that contains all the possibilities for $G_E(p)$ for any elliptic curve E/\mathbb{Q} and any prime p . It will turn out that this set is close enough to the truth for us to get a complete unconditional list of all possible degrees $[\mathbb{Q}(P) : \mathbb{Q}]$ for a large set of primes p .

Lemma 5.1 ([28, Theorem 5.1]). *Let E be an elliptic curve over a number field K and let p be a prime such that $G_E(p) = \text{GL}_2(\mathbb{F}_p)$. Then for a point P of order p , we have $[\mathbb{Q}(P) : \mathbb{Q}] = p^2 - 1$.*

Lemma 5.2. *Let E be an elliptic curve over a number field K and let p be a prime such that $G_E(p) = C_{ns}^+(p)$. Then for a point P of order p , we have $[\mathbb{Q}(P) : \mathbb{Q}] = p^2 - 1$.*

Proof. In [28, Lemma 7.5 (2)] it is claimed that $[\mathbb{Q}(P) : \mathbb{Q}] = p^2 - 1$ or $2(p^2 - 1)$. But the case $[\mathbb{Q}(P) : \mathbb{Q}] = 2(p^2 - 1)$ is not possible since it is impossible that $[\mathbb{Q}(P) : \mathbb{Q}] > (p^2 - 1)$ for any $P \in E[p]$ and any $G_E(p)$. This follows from the fact that $|G_E(p).v|$ can be at most $p^2 - 1$. The claim of the lemma follows. \square

Lemma 5.3 ([28, Lemma 7.5. (1)]). *Let E be an elliptic curve over a number field K and let p be a prime such that $G_E(p)$ is a conjugate of $C_s^+(p)$ in $\text{GL}_2(\mathbb{F}_p)$. Then for a point P of order p , we have $[\mathbb{Q}(P) : \mathbb{Q}]$ is either $(p - 1)^2$ or $2(p - 1)$; both cases occur.*

We will need a slightly finer description of the possibilities for $G_E(13)$ than the one supplied in Theorem 3.2.

Proposition 5.4. *Let E/\mathbb{Q} be a non-CM elliptic curve. Then $G_E(13)$ is conjugate in $\text{GL}_2(\mathbb{F}_{13})$ to one of the groups in Table 2, to $C_s^+(13)$, $C_{ns}^+(13)$, or one of the following groups:*

$$(7) \qquad 13Ns.2.1, \qquad 13Ns.5.2, \qquad 13Ns.5.1.4.$$

Proof. First recall that every group in Table 2 occurs. By Theorem 3.2(iii), the remaining possibilities are that $G_E(13)$ is a subgroup of $C_s^+(13)$, $C_{ns}^+(13)$, or 13S4. We say that G is an applicable subgroup of $\text{GL}_2(\mathbb{F}_{13})$ if $-I \in G$, $\det(G) = \mathbb{F}_{13}^\times$, and G contains an element with trace 0 and determinant -1 . Thanks to [37, Proposition 2.2] we have that if $G_E(13) \neq \text{GL}_2(\mathbb{F}_{13})$, then $\pm G_E(13)$ is an applicable subgroup of $\text{GL}_2(\mathbb{F}_{13})$. Using Magma we determine (see [14]) that $G_E(13)$ is conjugate in $\text{GL}_2(\mathbb{F}_{13})$ to $C_s^+(13)$, $C_{ns}^+(13)$, 13S4, or one of the following groups:

- 13Cs, 13Cs.1.1, 13Cs.1.11, 13Cs.1.3, 13Cs.1.4, 13Cs.1.6, 13Cs.1.8,
- 13Cs.12.1, 13Cs.12.3, 13Cs.12.4, 13Cs.3.1, 13Cs.3.4, 13Cs.4.1, 13Cs.5.1,
- 13Cs.5.4, 13Ns.2.1, 13Ns.5.1.4, 13Ns.5.2.

Now, it is not possible that $G_E(13)$ is conjugate to 13Cs or 13Cs.a.b, for any integers a, b , since then E would have two independent 13-isogenies over \mathbb{Q} . Therefore, E would be 13-isogenous over \mathbb{Q} to an elliptic curve with a 169-isogeny, whose existence contradicts Theorem 2.1. \square

Lemma 5.5. *Let E/\mathbb{Q} be an elliptic curve and let P be a point of order 13 in E . Then*

$$[\mathbb{Q}(P) : \mathbb{Q}] \in \{3, 4, 6, 12, 24, 39, 48, 52, 72, 78, 96, 144, 156, 168\}.$$

All the possible degrees occur.

Proof. The fact that all of the degrees above appear can be seen from Table 2 and from the degrees (see [14]) of $[\mathbb{Q}(P) : \mathbb{Q}]$ obtained from CM elliptic curves. The fact that no other degrees occur follows from Proposition 5.4 and from computing (see [14]) the lengths of orbits of all the possible images $G_E(p)$. \square

The next result gives the possible degrees $[\mathbb{Q}(P) : \mathbb{Q}]$ where P is a point of order a prime $p \neq 13$ on E such that $G_E(p)$ does not appear in Tables 1 and 2.

Theorem 5.6. *For an elliptic curve E/\mathbb{Q} with a given $G_E(p)$ in the left column, and varying through all $P \in E[p]$, the values of $[\mathbb{Q}(P) : \mathbb{Q}]$ obtained will be exactly the same as what appears in the right column.*

| | |
|----------------------|--------------------------------|
| $G_E(p)$ | $[\mathbb{Q}(P) : \mathbb{Q}]$ |
| $GL_2(\mathbb{F}_p)$ | $p^2 - 1$ |
| $C_{ns}^+(p)$ | $p^2 - 1$ |
| $C_s^+(p)$ | $2(p - 1), (p - 1)^2$ |

| | |
|-------------|--------------------------------|
| $G_E(p)$ | $[\mathbb{Q}(P) : \mathbb{Q}]$ |
| $G_{00}(p)$ | $p - 1, (p - 1)p$ |
| $G_{01}(p)$ | $p - 1, (p - 1)p/2$ |
| $G_{10}(p)$ | $(p - 1)/2, (p - 1)p$ |

$p \in \{3, 7, 11, 19, 43, 67, 163\}$

| | |
|-----------------------------------|---------------------------------------|
| $G_E(p)$ | $[\mathbb{Q}(P) : \mathbb{Q}]$ |
| $G^3(p)$ $p \equiv 1 \pmod{3}$ | $2(p - 1), (p - 1)^2/3, 2(p - 1)^2/3$ |
| $G_0(p)$ $p \equiv 2 \pmod{3}$ | $(p^2 - 1)/3, 2(p^2 - 1)/3$ |

Remark 5.7. The groups $G_{00}(p)$, $G_{01}(p)$, $G_{10}(p)$, $G^3(p)$, and $G_0(p)$ are as defined in Theorems 3.2 and 3.6.

Proof. The first table follows from Lemmas 5.1, 5.2, and 5.3. The second table follows from a direct computation (see [14]), as one has to consider only a small number of finite group actions.

Now let $G_E(p) = G^3(p)$. Then it is enough to consider the lengths orbits of the generators of the $p + 1$ subgroups of \mathbb{F}_p^2 . Equivalently, it is enough to know the size of the stabilizers of each of these elements. The stabilizer of $(1, 0)$ is by [28, Lemma 6.6 (1)] the subgroup of $G^3(p)$ given by

$$H_{(1,0)} = \{D(1, b) : b \in \mathbb{F}_p^\times\}.$$

Obviously, the stabilizer is of size $(p - 1)/3$, hence the size of the orbit of $(1, 0)$ is $\frac{|G^3(p)|}{(p-1)/3} = 2(p - 1)$. Using [28, Lemma 6.6] and the same argumentation, we conclude that the length of the orbit of $(0, 1)$ is also $2(p - 1)$. The stabilizer of any element of the form $(1, x)$ for $x \in \mathbb{F}_p^\times$ in $C_s^+(p)$ is by [28, Lemma 6.6 (3)]

$$H_{(1,x)} = \{D(1, 1), D(x, x^{-1})\}.$$

Thus the stabilizer of $(1, x)$ in $G^3(p)$ is $H \cap G^3(p)$; this set obviously contains either 1 or 2 elements. It contains 2 elements, by the definition of $G^3(p)$, if and only if x^2 is a cube in \mathbb{F}_p^\times . Obviously there exist both x 's that satisfy this condition (take for example x to be a cube itself) and x that do not. Hence there exist vectors $(1, x)$ which are in an orbit of length $\frac{(p-1)^2}{3}$ and those that are in an orbit of length $\frac{2(p-1)^2}{3}$. In this case \mathbb{F}_p^2 decomposes into orbits of lengths $1, \frac{(p-1)^2}{3}, \frac{2(p-1)^2}{3}$.

Now consider the case $G_E(p) = G_0(p)$. It follows from [28, Lemma 7.4], that the stabilizer of each element is of order 1 or 2. If there exists an element whose stabilizer is of order 1, then \mathbb{F}_p^2 decomposes into orbits of length $1, \frac{p^2-1}{3}, \frac{2(p^2-1)}{3}$,

and if there does not exist such an element, then \mathbb{F}_p^2 decomposes in orbits of length $1, \frac{p^2-1}{3}, \frac{p^2-1}{3}, \frac{p^2-1}{3}$ under the action of $G_E(p)$.

By [28, Lemma 7.4], the elements of $C_{ns}^+(p)$ that fix any non-trivial vector (or equivalently have an eigenvalue 1) are the identity and the matrices $J \cdot M_\epsilon(a, b)$ such that $a^2 - \epsilon b^2 = 1$, or equivalently those such that $\det J \cdot M_\epsilon(a, b) = -1$. It is easy to see that there are $p + 1$ such matrices in $C_{ns}^+(p)$, each fixing each element in a different subgroup of $(\mathbb{F}_p)^2$.

We now study the number of such elements in $G_0(p)$. Clearly, none of them are of the form $M_\epsilon(a, b)^3$, so all of them must be of the form $J \cdot M_\epsilon(a, b)^3$.

Now, $J \cdot M_\epsilon(a, b)^3$ has determinant -1 if and only if $M_\epsilon(a, b)^3$ has determinant 1. Since 1 is the only third root of unity in \mathbb{F}_p , it follows that $\det(M_\epsilon(a, b)^3) = 1$ if and only if $\det M_\epsilon(a, b) = 1$. The subgroup \mathcal{M} of $C_{ns}(p)$ of matrices $M_\epsilon(a, b)$ of determinant 1 is the kernel of the determinant map $\det : C_{ns}(p) \rightarrow \mathbb{F}_p^\times$, which is surjective. Hence, by the first isomorphism theorem, $\#\mathcal{M} = p + 1$. As $\#\mathcal{M}$ is divisible by 3, it follows that $\mathcal{M}^3 = \{M_\epsilon(a, b)^3 : (a, b) \in (\mathbb{F}_p)^2, (a, b) \neq (0, 0)\}$ has cardinality $\frac{p+1}{3}$, as claimed.

Hence there are $\frac{p+1}{3}$ non-identity matrices that each fix every element in exactly one subgroup of $(\mathbb{F}_p)^2$ of order p . As there are $p + 1$ subgroups of $(\mathbb{F}_p)^2$ of order p , it follows that there will be $\frac{2(p+1)}{3}$ subgroups such that each of their non-trivial elements has trivial stabilizer, proving the claim. □

We now state the main result of this section.

Theorem 5.8. *Let E/\mathbb{Q} be an elliptic curve, let p be a prime, and let P be a point of order p in E . Then all of the cases in the table below occur for $p \leq 13$ or $p = 37$, and they are the only ones possible. The degrees in the table and lists below with an asterisk occur only when E has CM.*

| p | $[\mathbb{Q}(P) : \mathbb{Q}]$ |
|-----|---|
| 2 | 1, 2, 3 |
| 3 | 1, 2, 3, 4, 6, 8 |
| 5 | 1, 2, 4, 5, 8, 10, 16, 20, 24 |
| 7 | 1, 2, 3, 6, 7, 9, 12, 14, 18, 21, 24*, 36, 42, 48 |
| 11 | 5, 10, 20*, 40*, 55, 80*, 100*, 110, 120 |
| 13 | 3, 4, 6, 12, 24*, 39, 48*, 52, 72, 78, 96, 144*, 156, 168 |
| 37 | 12, 36, 72*, 444, 1296*, 1332, 1368 |

For all other p , the possibilities for $[\mathbb{Q}(P) : \mathbb{Q}]$ are as is given below. The degrees in equations (10)-(12) occur only for CM elliptic curves E/\mathbb{Q} . Furthermore the degrees in equations (12) occur only for elliptic curves with j -invariant 0. If Conjecture 3.5 is true, then the degrees in equations (13) also occur only for elliptic curves with j -invariant 0. Then

- (8) $p^2 - 1$ for all p ,
- (9) $8, 16, 32^*, 136, 256^*, 272, 288$ for $p = 17$,
- (10) $(p - 1)/2, p - 1, p(p - 1)/2, p(p - 1)$ if $p \in \{19, 43, 67, 163\}$,

$$(11) \quad 2(p-1), \quad (p-1)^2 \quad \text{if } p \equiv 1 \pmod{3} \text{ or } \left(\frac{-D}{p}\right) = 1$$

for any $D \in \mathcal{CM}$,

$$(12) \quad (p-1)^2/3, \quad 2(p-1)^2/3 \quad \text{if } p \equiv 4, 7 \pmod{9},$$

$$(13) \quad (p^2-1)/3, \quad 2(p^2-1)/3 \quad \text{if } p \equiv 2, 5 \pmod{9},$$

where $\mathcal{CM} = \{1, 2, 7, 11, 19, 43, 67, 163\}$.

Apart from the cases above that have been proven to appear, the only other options that might be possible are:

$$(14) \quad (p^2-1)/3, \quad 2(p^2-1)/3 \quad \text{if } p \equiv 8 \pmod{9}.$$

Proof. The theorem follows directly from the results of this section, combined with Theorems 3.2 and 3.6, and from direct calculations (see [14]) of lengths of orbits of fixed finite groups. □

Remark 5.9. Note that 3 would be redundant in the set \mathcal{CM} , although there exists an elliptic curve with CM by an order of an imaginary quadratic field of discriminant -3 , as $p \equiv 1 \pmod{3}$ immediately implies $\left(\frac{-D}{p}\right) = -1$. That is why 3 has been left out from the set \mathcal{CM} .

Remark 5.10. If Conjecture 3.5 is true, then the cases (14) do not occur.

6. POINTS OF PRIME ORDER ON ELLIPTIC CURVES E/\mathbb{Q} OVER NUMBER FIELDS OF DEGREE d

Lozano-Robledo [28] denotes by $S_{\mathbb{Q}}(d)$ the set of primes p for which there exists a number field K of degree $\leq d$ and an elliptic curve E/\mathbb{Q} such that the p divides the order of $E(K)_{\text{tors}}$. He determined the sets $S_{\mathbb{Q}}(d)$ for $d \leq 42$. To more easily state his results, denote by $S_{\mathbb{Q}}^*(d) = S_{\mathbb{Q}}(d) \setminus S_{\mathbb{Q}}(d-1)$; then the table below shows the necessary data to recover $S_{\mathbb{Q}}(d)$ for $d \leq 42$:

| | | | | | | | | | |
|-----------------------|------------|----|----|----|----|----|----|----|-------------------|
| d | 1 | 3 | 5 | 8 | 9 | 12 | 21 | 33 | other $d \leq 42$ |
| $S_{\mathbb{Q}}^*(d)$ | 2, 3, 5, 7 | 13 | 11 | 17 | 19 | 37 | 43 | 67 | \emptyset |

In this section, we obtain results about a set that will give us more information: $R_{\mathbb{Q}}(d)$. Recall from the introduction that we denote by $R_{\mathbb{Q}}(d)$ the set of primes p for which there exists a number field K of degree exactly d and an elliptic curve E/\mathbb{Q} such that the p divides the order of $E(K)_{\text{tors}}$. Obviously, knowing $R_{\mathbb{Q}}(d)$ immediately gives $S_{\mathbb{Q}}(d)$, but not vice versa. We obtain our results as more or less direct consequences of Theorem 5.8.

The first obvious observation is that for all positive integers n, d , if $p \in R_{\mathbb{Q}}(d)$, then $p \in R_{\mathbb{Q}}(nd)$. Obviously, we have $2, 3, 5, 7 \in R_{\mathbb{Q}}(d)$ for all positive integers d .

The following corollary summarizes the results that we obtain.

Corollary 6.1. *Let $\mathcal{CM} = \{1, 2, 7, 11, 19, 43, 67, 163\}$. The following holds:*

- (i) $11 \in R_{\mathbb{Q}}(d)$ if and only if $5|d$.
- (ii) $13 \in R_{\mathbb{Q}}(d)$ if and only if $3|d$ or $4|d$.
- (iii) $17 \in R_{\mathbb{Q}}(d)$ if and only if $8|d$.
- (iv) $37 \in R_{\mathbb{Q}}(d)$ if and only if $12|d$.
- (v) For $p = 19, 43, 67, 163$, it holds that $p \in R_{\mathbb{Q}}(d)$ if and only if $\frac{p-1}{2}|d$.

- (vi) For $p \geq 23$, $p \neq 37, 43, 67, 163$:
 - (a) If $p \equiv 1 \pmod{3}$, then it holds that $p \in R_{\mathbb{Q}}(d)$ if and only if $2(p-1)|d$.
 - (b) If $p \equiv 2 \pmod{3}$ and $\left(\frac{-D}{p}\right) = 1$ for at least one of the values $D \in \mathcal{CM}$, then $p \in R_{\mathbb{Q}}(d)$ if and only if $2(p-1)|d$.
 - (c) If $p \equiv 2, 5 \pmod{9}$ and $\left(\frac{-D}{p}\right) = -1$ for all the values $D \in \mathcal{CM}$, then $p \in R_{\mathbb{Q}}(d)$ if and only if $(p^2-1)/3|d$.
 - (d) Let $p \equiv 8 \pmod{9}$ and $\left(\frac{-D}{p}\right) = -1$ for all the values $D \in \mathcal{CM}$, then if $p \in R_{\mathbb{Q}}(d)$, then it follows that $\frac{p^2-1}{3} | d$. On the other hand if $p^2-1|d$, then it follows that $p \in R_{\mathbb{Q}}(d)$.

Proof. This follows directly from Theorem 5.8 and direct computations. □

Remark 6.2. Note that the only case where we do not have one necessary and sufficient condition for p to be in $R_{\mathbb{Q}}(d)$ is (vi)(d). This is because of the possibility that $G_E(p) = G_0(p)$ in that case. If Conjecture 3.5 is true, then $G_E(p) = G_0(p)$ will be impossible for all $p > 13$, and we will have $p \in R_{\mathbb{Q}}(d)$ if and only if $p^2-1|d$ in this case.

Of course, it is a natural question to ask what is the density of primes satisfying the assumptions of (vi)(d). Notice that $\left(\frac{-4}{p}\right) = \left(\frac{-1}{p}\right) = -1$ is true if and only if $p \equiv 3 \pmod{4}$. The condition $\left(\frac{-2}{p}\right) = -1$ is then equivalent to $\left(\frac{2}{p}\right) = 1$, which implies that $p \equiv 7 \pmod{8}$.

Now, by quadratic reciprocity,

$$\left(\frac{-D}{p}\right) = \left(\frac{-1}{p}\right) \left(\frac{D}{p}\right) = (-1) \cdot (-1) \cdot \left(\frac{p}{D}\right) = \left(\frac{p}{D}\right)$$

for all $D \in \mathcal{CM} - \{1, 2\}$. For each of these D , the condition $\left(\frac{p}{D}\right) = -1$ is satisfied by half of the residue classes modulo D . Combining, by the Chinese remainder theorem, all these conditions together with $p \equiv 7 \pmod{8}$ and $p \equiv 8 \pmod{9}$ we obtain that the assumptions of (vi) (d) are satisfied by $\frac{1}{4} \cdot \frac{1}{6} \cdot \left(\frac{1}{2}\right)^6 = \frac{1}{1536}$ of the residue classes modulo $8 \cdot 9 \cdot 7 \cdot 11 \cdot 19 \cdot 43 \cdot 67 \cdot 163$. As the primes are evenly distributed in residue classes, we conclude that a set of natural density $\frac{1}{1536}$ of primes satisfies the assumptions of (vi)(d).

Remark 6.3. Let p be a prime and denote by \mathcal{D}_p the minimal subset of the possible degrees $[\mathbb{Q}(P) : \mathbb{Q}]$ for any point P of order p in any elliptic curve E/\mathbb{Q} , such that $[\mathbb{Q}(P) : \mathbb{Q}]$ is divisible by some $d \in \mathcal{D}_p$. Then

$$R_{\mathbb{Q}}(d) = \{p \text{ prime} : \exists a \in \mathcal{D}_p \text{ such that } a|d\}.$$

Corollary 6.4. *Let $p = 2$ or $p > 5$ be a prime. Then $R_{\mathbb{Q}}(p) = R_{\mathbb{Q}}(1) = \{2, 3, 5, 7\}$.*

In this context we denote by

$$R_{\mathbb{Q}}^*(d) = R_{\mathbb{Q}}(d) \setminus \bigcup_{\substack{d'|d \\ d' \neq d}} R_{\mathbb{Q}}(d').$$

The table below shows the data (see [14]) needed to recover $R_{\mathbb{Q}}(d)$ for $d \leq 100$:

| | | | | | | | | | | | | | | | |
|-----------------------|------------|------|----|----|----|----|----|----|----|----|----|----|-----|----|-----------------------|
| d | 1 | 3, 4 | 5 | 8 | 9 | 12 | 21 | 33 | 44 | 56 | 60 | 80 | 81 | 92 | other $d \leq 100$ |
| $R_{\mathbb{Q}}^*(d)$ | 2, 3, 5, 7 | 13 | 11 | 17 | 19 | 37 | 43 | 67 | 23 | 29 | 31 | 41 | 163 | 47 | \emptyset |

Note that the smallest prime that satisfies the condition of Corollary 6.1(vi)(d) is $p = 3167$. Therefore, the smallest d where we do not have an unconditional determination of $R_{\mathbb{Q}}(d)$ is $d = 3343296$.

7. TORSION GROUPS OF ELLIPTIC CURVES E/\mathbb{Q}
OVER NUMBER FIELDS OF DEGREE d

In this section we want to say things about the possibilities for the whole torsion group of elliptic curves E/\mathbb{Q} over number fields, not just about points of prime degree, as in the previous section.

We write $E(K)[p^\infty]$ to denote the p -primary torsion subgroup of $E(K)_{\text{tors}}$, that is, the p -Sylow subgroup of $E(K)$.

We will prove results about elliptic curves in prime degree fields. We first need to study the case of degree 7 separately.

Recall from the introduction that $\Phi_{\mathbb{Q}}(d) \subseteq \Phi(d)$ is the set of possible isomorphism classes of groups $E(K)_{\text{tors}}$, where K runs through all number fields K of degree d and E runs through all elliptic curves defined over \mathbb{Q} .

Proposition 7.1. $\Phi_{\mathbb{Q}}(7) = \Phi(1)$.

Proof. Throughout the proof let E/\mathbb{Q} be an elliptic curve and let K be a number field of degree 7. First suppose that E has CM. We can see from [6, Theorem 1.4] (or [8, §4.7]) that in the CM case, no subgroups occur in degree 7 which do not occur over \mathbb{Q} .

Now assume E/\mathbb{Q} is non-CM. First we show that $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ for $p \neq 7$ prime. For $p \leq 11$, it follows that $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ from Table 1 in the appendix (note that by [37], this list of mod p Galois images for $p \leq 11$ is complete). For $p \geq 13$ it follows directly from Corollary 6.1.

It remains to deal with the case $p = 7$. From Table 1 we see that $E(\mathbb{Q})[7] = E(K)[7]$ unless $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3. We claim that if $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3, then $E(\mathbb{Q})[p] = \{\mathcal{O}\}$ for all $p \neq 7$. Indeed, such an elliptic curve would have a $7p$ -isogeny, and this is possible only for $p = 2$ and $p = 3$, and for only finitely many elliptic curves, up to $\overline{\mathbb{Q}}$ -isomorphism. Note that if $j_E \notin \{0, 1728\}$ and E' is a quadratic twist of E , then $\pm G_E(n) = \pm G_{E'}(n)$ for any integer $n \geq 3$. We explicitly check (see [14]) that no elliptic curve with a 14-isogeny or a 21-isogeny has $G_E(7)$ conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3.

It remains to prove that $E(K)_{\text{tors}}$ contains no points of order 49. Looking at Table 1 we observe that a necessary condition for $E(K)_{\text{tors}}$ to contain a point of order 7 is that $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.1 or 7B.1.3. Assume that there exists a number field K of degree 7 such that $E(K)[49] = \langle P \rangle \simeq \mathcal{C}_{49}$. Then $G_E(49)$ satisfies:

$$G_E(49) \equiv G_E(7) \pmod{7} \quad \text{and} \quad [G_E(49) : H_P] = 7,$$

where H_P is the image in $\text{GL}_2(\mathbb{Z}/49\mathbb{Z})$ of the subgroup $\mathcal{H}_P \subseteq \text{Gal}(\mathbb{Q}(E[49])/\mathbb{Q})$ such that $\mathbb{Q}(P) = \mathbb{Q}(E[49])^{\mathcal{H}_P}$.

Note that in general we do not have an explicit description of possible images $G_E(49)$, but using **Magma** we can run through all subgroups of $\text{GL}_2(\mathbb{Z}/49\mathbb{Z})$ that reduce to $G_E(7)$.

First assume that $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3. We check (see [14]) using **Magma** that for any subgroup G of $\text{GL}_2(\mathbb{Z}/49\mathbb{Z})$ satisfying $G \equiv H \pmod{7}$ for some conjugate H of 7B.1.3 in $\text{GL}_2(\mathbb{Z}/7\mathbb{Z})$, and for any $v \in (\mathbb{Z}/49\mathbb{Z})^2$ of order

49, we have $[G : G_v] \neq 7$, where G_v is the stabilizer of v by the action of G on $(\mathbb{Z}/49\mathbb{Z})^2$. Therefore for any point $P \in E(\overline{\mathbb{Q}})$ of order 49 we have $[G_E(49) : H_P] \neq 7$. In particular this proves that if $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3, then $E(K)_{\text{tors}}$ contains no points of order 49.

Now assume that $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.1. Using a similar procedure as the one used before, we search (see [14]) for subgroups G of $\text{GL}_2(\mathbb{Z}/49\mathbb{Z})$ satisfying $G \equiv H \pmod{7}$ for some subgroup H conjugate to 7B.1.1 in $\text{GL}_2(\mathbb{Z}/7\mathbb{Z})$, and satisfying $[G : G_v] = 7$ for some $v \in (\mathbb{Z}/49\mathbb{Z})^2$ of order 49. There exist such groups G , but in all the cases we obtain that G_v is a normal subgroup of G and $G/G_v \simeq \mathcal{C}_7$. Therefore we have deduced that if E/\mathbb{Q} is an elliptic curve such that $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.1 and there exists a number field K of degree 7 with a K -rational point of order 49, then K is Galois. But this would imply that E has a rational 49-isogeny, and that is impossible by Theorem 2.1.

This finishes the proof. □

Theorem 7.2. *Let B be a positive integer. Let E/\mathbb{Q} be an elliptic curve and let K/\mathbb{Q} be a number field of degree d , where the smallest prime divisor of d is $\geq B$. Then*

- (i) *If $B \geq 11$, then $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ for all primes p . In particular, $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$.*
- (ii) *If $B \geq 7$, then $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ for all primes $p \neq 7$.*
- (iii) *If $B \geq 5$, then $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ for all primes $p \neq 5, 7, 11$.*
- (iv) *If $B > 2$, then $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$ for all primes $p \neq 2, 3, 5, 7, 11, 13, 19, 43, 67, 163$.*

Proof. We will prove only the case (i)—the cases (ii), (iii), and (iv) are proved using the exact same argumentation. Suppose first that $P \in E(\overline{\mathbb{Q}})$ is a point of order $p \leq 13$. Then directly (and obviously) from Theorem 5.6 we see that $[\mathbb{Q}(P) : \mathbb{Q}]$ has to be divisible by some element from the set $\{2, 3, 5, 7\}$.

For $p \geq 17$, by looking at the possibilities listed in Theorem 5.6 we have that $[\mathbb{Q}(P) : \mathbb{Q}]$ always attains only obviously even values, except in the cases

$$[\mathbb{Q}(P) : \mathbb{Q}] = \frac{p-1}{2} \quad \text{and} \quad [\mathbb{Q}(P) : \mathbb{Q}] = \frac{p(p-1)}{2},$$

which are possible only for $p \in \{3, 7, 11, 19, 43, 67, 163\}$. Since the cases $p \in \{3, 7, 11\}$ have already been dealt with, it remains to look at only the cases $p \in \{19, 43, 67, 163\}$. Since in all of these cases we have $p \equiv 1 \pmod{3}$, it follows that $\frac{p-1}{2}$ will be divisible by 3. We conclude that $E(K)[p] = E(\mathbb{Q})[p]$ for all primes p if K satisfies the assumptions of the theorem. Now it follows from Proposition 4.6 that $E(K)[p^\infty] = E(\mathbb{Q})[p^\infty]$. □

We immediately obtain the following result.

Corollary 7.3. *Let d be a positive integer such that the smallest prime factor of d is ≥ 11 . Then $\Phi_{\mathbb{Q}}(d) = \Phi(1)$.*

Remark 7.4. One could show, using arguments as in the proof of Proposition 4.6, that for any $p \in \{2, 3, 5, 7\}$, the set

$$\bigcup_{n=1}^{\infty} \Phi_{\mathbb{Q}}(p^n)$$

will be infinite, containing \mathcal{C}_{p^k} for all positive integers k .

So, in a sense, Corollary 7.3 is best possible.

Remark 7.5. From Corollary 7.3 and the previous remark it follows that the d such that $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$ for all elliptic curves E/\mathbb{Q} and all number fields K of degree d are exactly those such that the smallest prime divisor of d is ≥ 11 . These are d of the form $d = 210k + x$, where k is a non-negative integer and $1 \leq x < 210$ is an integer coprime to 210. Thus it follows that exactly $\frac{\phi(210)}{210} = \frac{8}{35}$ of all integers (when ordered by size) satisfy this property.

Our methods give results not only about the torsion in finite extensions, but also in infinite extensions of \mathbb{Q} .

Corollary 7.6. *Let $p \geq 11$ be a prime and let K be the \mathbb{Z}_p -extension of \mathbb{Q} . Then $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$.*

Proof. Let ℓ be a prime. First suppose that $E(\mathbb{Q})[\ell] = \{\mathcal{O}\}$. Then we have $E(F)[\ell] = \{\mathcal{O}\} = E(\mathbb{Q})_{\text{tors}}$ by Theorem 7.2 for all number fields F contained in K .

Now suppose that $E(\mathbb{Q})[\ell] \neq \{\mathcal{O}\}$. Then it follows that $\ell \leq 7$, and hence $p \neq \ell$. If $\ell = 2$, then it is obvious that $E(\mathbb{Q})[2] = E(K)[2]$. If $3 \leq \ell \leq 7$, then $E(\mathbb{Q})[\ell] = E(K)[\ell]$, since K does not contain ℓ th roots of unity.

By Proposition 4.6, we now have $E(\mathbb{Q})[\ell^\infty] = E(F)[\ell^\infty]$ for all number fields F contained in K , from which we again conclude that $E(\mathbb{Q})[\ell^\infty] = E(K)[\ell^\infty]$. \square

7.1. Torsion growth in degree 7 extensions. In this section we consider how the torsion grows in prime degree extensions in the following sense: for each possible $E(\mathbb{Q})_{\text{tors}}$, what can $E(K)_{\text{tors}}$ possibly be, for K/\mathbb{Q} an extension of prime degree d . There is no torsion growth in degree $p \geq 11$ by Theorem 7.2 and the torsion growth for $p \leq 5$ has been completely described in [16] for $p = 2$, in [15] for $p = 3$, and in [11] for $p = 5$. Thus it remains to consider the case $p = 7$. We prove the following.

Proposition 7.7. *Let E/\mathbb{Q} be an elliptic curve and let K be a number field of degree 7.*

- (i) *If $E(\mathbb{Q})_{\text{tors}} \not\cong \mathcal{C}_1$, then $E(\mathbb{Q})_{\text{tors}} = E(K)_{\text{tors}}$.*
- (ii) *If $E(\mathbb{Q})_{\text{tors}} \cong \mathcal{C}_1$, then $E(K)_{\text{tors}} \cong \mathcal{C}_1$ or \mathcal{C}_7 . Furthermore, if $E(\mathbb{Q})_{\text{tors}} \cong \mathcal{C}_1$ and $E(K)_{\text{tors}} \cong \mathcal{C}_7$, then K is the unique degree 7 number field with this property and E is isomorphic to the elliptic curve*

$$E_t : y^2 = x^3 + 27(t^2 - t + 1)(t^6 + 229t^5 + 270t^4 - 1695t^3 + 1430t^2 - 235t + 1)x + 54(t^{12} - 522t^{11} - 8955t^{10} + 37950t^9 - 70998t^8 + 131562t^7 - 253239t^6 + 316290t^5 - 218058t^4 + 80090t^3 - 14631t^2 + 510t + 1)$$

for some value $t \in \mathbb{Q}$.

Proof. Let E/\mathbb{Q} be an elliptic curve. Clark, Corn, Rice, and Stankewicz [8, §4.7] have proved that if E has CM, then $E(K)_{\text{tors}}$ has torsion points only of order 1, 2, 3, 4, or 6. Now from Lemma 3.1, it follows that $E(K)[n] = E(\mathbb{Q})[n]$ for $n = 2, 3, 4, 6$. It follows that $E(K)_{\text{tors}} = E(\mathbb{Q})_{\text{tors}}$.

Then we can assume that E/\mathbb{Q} is non-CM. In the proof of Proposition 7.1 we proved that if p is a prime and K/\mathbb{Q} is a number field of degree 7, then $E(\mathbb{Q})[p^\infty] = E(K)[p^\infty]$, unless $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to **7B.1.3**, in which case $E(\mathbb{Q})[p] = \{\mathcal{O}\}$ for all $p \neq 7$. Then suppose that $G_E(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to **7B.1.3**. Zywinia [37, Theorem 1.5 (iv)] proved that, under these assumptions, there exists a $t_0 \in \mathbb{Q}$ such that E is \mathbb{Q} -isomorphic to E_{t_0} .

Let $\psi_7(x)$ be the 7-division polynomial of E_t and let $\psi_7(x) = f_1(x)f_2(x)f_3(x)f_4(x)$ be its irreducible factorization (over $\mathbb{Q}(t)$), where $f_1(x)$ is of degree 3 and $f_i(x)$, for $i = 2, 3, 4$, are all of degree 7. For all specializations E_{t_0} , $t_0 \in \mathbb{Q}$, we have that $G_{E_{t_0}}(7)$ is conjugate in $\text{GL}_2(\mathbb{F}_7)$ to 7B.1.3, as it cannot be any smaller, so the 7-division polynomial of E_{t_0} will factor over \mathbb{Q} in the same manner as the 7-division polynomial of E_t factors over $\mathbb{Q}(t)$. We compute (see [14]) that the roots of $f_i(x)$, for $i = 2, 3, 4$, define the same degree 7 extensions of $\mathbb{Q}(t)$, up to isomorphism. We obtain that E_t has torsion \mathcal{C}_7 over this extension, completing the proof. \square

Remark 7.8. Let E/\mathbb{Q} be an elliptic curve and let p be a prime. If the possible degrees of points of order p are p and $p - 1$, then $\mathbb{Q}(E[p]) = K\mathbb{Q}(\zeta_p)$ satisfies $[\mathbb{Q}(E[p]) : \mathbb{Q}] = p(p-1)$ and there exists $P \in E[p]$ such that $K = \mathbb{Q}(P)$ is a Kummer extension, i.e., there exists $s \in \mathbb{Z}$, a non- p th-power, such that $K = \mathbb{Q}(\sqrt[p]{s})$. This is the case when $G_E(p)$ is conjugate in $\text{GL}_2(\mathbb{F}_p)$ to 3B.1.2, 5B.1.2 or 7B.1.3.

8. TORSION OF ELLIPTIC CURVES E/\mathbb{Q} OVER QUARTIC NUMBER FIELDS

In this section we solve the problem of determining the set of $\Phi_{\mathbb{Q}}(4)$. The sets $\Phi_{\mathbb{Q}}(2)$ and $\Phi_{\mathbb{Q}}(3)$ have been determined by the second author [32], $\Phi_{\mathbb{Q}}(5)$ by the first author [11].

Let G be a finite group; define $\Phi_{\mathbb{Q}}^G(n)$ to be the set of isomorphism classes of groups $E(K)_{\text{tors}}$, where E is an elliptic curve defined over \mathbb{Q} and $\text{Gal}(\widehat{K}/\mathbb{Q}) \simeq G$, where \widehat{K} denotes the normal closure of K over \mathbb{Q} . The first steps taken towards determining $\Phi_{\mathbb{Q}}(4)$ have been done by Chou [7], who determined $\Phi_{\mathbb{Q}}^{\mathcal{C}_4}(4)$ and $\Phi_{\mathbb{Q}}^{S_4}(4)$, i.e., he found the possible torsion groups over Galois quartic number fields of elliptic curves defined over \mathbb{Q} .

The strategy used both in [7] and [32] (for determining $\Phi_{\mathbb{Q}}(3)$) is to view how $\text{Gal}(\widehat{K}/\mathbb{Q})$ acts on $E(K)_{\text{tors}}$ and to conclude that this forces certain properties on E over \mathbb{Q} . For certain choices of $E(K)_{\text{tors}}$, one would get properties of E over \mathbb{Q} that would violate known results about elliptic curves over \mathbb{Q} , thus ruling out the possibility of that particular torsion group $E(K)_{\text{tors}}$. This approach works well when K is Galois over \mathbb{Q} , as in [7], or “not far from Galois” in [32] where $\text{Gal}(\widehat{K}/\mathbb{Q})$ was either \mathcal{C}_3 or S_3 (even in this case, S_3 posed some problems). We make use of the machinery developed in Section 4 to show obstructions for possible growth from $E(\mathbb{Q})_{\text{tors}}$ to $E(K)_{\text{tors}}$ when K is “far away from being Galois” (meaning informally that $[\widehat{K} : K]$ is relatively large).

The results of Section 4 immediately tell us that (see Corollary 4.2)

$$\Phi_{\mathbb{Q}}^{A_4}(4) = \Phi(1).$$

In some instances in the remainder of the paper we will say that a quartic field K has Galois group D_4 , A_4 , or S_4 , which will always mean that its normal closure \widehat{K} satisfies $\text{Gal}(\widehat{K}/\mathbb{Q}) \simeq D_4$, A_4 , or S_4 , respectively.

We will need to look at the case of the possible 3-torsion growth in a quartic number field with Galois group S_4 separately, as this case is a bit more delicate.

Proposition 8.1. *Let E/\mathbb{Q} be an elliptic curve and let K be a quartic number field with Galois group isomorphic to S_4 . Then $E(K)[3] = E(\mathbb{Q})[3]$.*

Proof. Suppose that $E(\mathbb{Q})[3] \subsetneq E(K)[3]$. Then by Theorem 4.3, we have that S_4 is isomorphic to a quotient of a subgroup of $G_E(3)$. We deduce that $G_E(3) = \text{GL}_2(\mathbb{F}_3)$,

since $\#S_4 = 24$, $\#GL_2(\mathbb{F}_3) = 48$, and $GL_2(\mathbb{F}_3)$ has (see [14]) no subgroups isomorphic to S_4 . But then the number field of smallest degree over which there exists a point P of order 3 satisfies

$$\text{Gal}(\mathbb{Q}(E[3])/\mathbb{Q}(P)) \simeq \left\{ \begin{pmatrix} 1 & c \\ 0 & d \end{pmatrix} : c \in \mathbb{F}_3, d \in \mathbb{F}_3^\times \right\},$$

so by Galois theory is a number field of degree 8 over \mathbb{Q} . □

Remark 8.2. We should note that if $G_E(3) = GL_2(\mathbb{F}_3)$, as in the proof above, then $\mathbb{Q}(E[3])$ in fact contains a Galois extension K of \mathbb{Q} with Galois group isomorphic to S_4 . This will be the field generated by all the x -coordinates of $E[3]$. However, it will hold that $E(K)[3] = \{\mathcal{O}\}$.

Now we obtain the following result.

Corollary 8.3. *Let $p \geq 3$ be a prime, let n be a positive integer, and let K be a quartic number field with Galois group isomorphic to A_4 or S_4 . If an elliptic curve E/\mathbb{Q} has no points of order p^n in $E(\mathbb{Q})$, then it has no points of order p^n in $E(K)$.*

Proof. For $n \geq 2$, the result follows from Theorem 4.5. Suppose from now on that $n = 1$. If $\text{Gal}(\widehat{K}/\mathbb{Q}) \simeq S_4$ and $p = 3$ (resp., $p \geq 5$) the results follow from Proposition 8.1 (resp., Theorem 4.3). If $\text{Gal}(\widehat{K}/\mathbb{Q}) \simeq A_4$ the result follows from Theorem 4.3 for any $p \geq 3$. □

As a corollary of Proposition 4.8, we can now obtain the following results about subsets of $\Phi_{\mathbb{Q}}(4)$.

Corollary 8.4. $\Phi_{\mathbb{Q}}^{S_4}(4) = \Phi_{\mathbb{Q}}^{A_4}(4) = \Phi(1)$.

Proof. Suppose K is a quartic number field with Galois group A_4 or S_4 .

Let E/\mathbb{Q} be an elliptic curve. Because of Corollary 4.2 and Proposition 8.1, we have $E(\mathbb{Q})[p] = E(K)[p]$ for all $p \geq 3$. Due to Corollary 8.3, we have that there are no points of order p^n over K for $p \geq 3$ and $n \geq 2$.

For $p = 2$, we note that $\mathbb{Q}(E[2])$ is either \mathbb{Q} , a quadratic field, cyclic cubic, or an S_3 extension of \mathbb{Q} and the intersection of K with any of these fields is \mathbb{Q} , so $E(\mathbb{Q})[2] = E(K)[2]$. Furthermore, by Proposition 4.8, if $E(K)[2^n] \supsetneq E(\mathbb{Q})[2^n]$, for $n \geq 2$, then the Galois group of K is isomorphic to a subgroup of D_4 , which is a contradiction. □

Thus to determine $\Phi_{\mathbb{Q}}(4)$ completely (see Corollary 8.7), it remains to just determine $\Phi_{\mathbb{Q}}^{D_4}(4)$.

8.1. Determining $\Phi_{\mathbb{Q}}^{D_4}(4)$. In this section we determine $\Phi_{\mathbb{Q}}^{D_4}(4)$. In particular, we prove the following theorem.

Theorem 8.5. *The set $\Phi_{\mathbb{Q}}^{D_4}(4)$ is given by*

$$\begin{aligned} \Phi_{\mathbb{Q}}^{D_4}(4) = & \{ \mathcal{C}_n : n = 1, \dots, 10, 12, 15, 16, 20, 24 \} \cup \{ \mathcal{C}_2 \times \mathcal{C}_{2m} : m = 1, \dots, 6 \} \\ & \cup \{ \mathcal{C}_3 \times \mathcal{C}_{3m} : m = 1, 2 \} \cup \{ \mathcal{C}_4 \times \mathcal{C}_{4m} : m = 1, 2 \}. \end{aligned}$$

Remark 8.6. $\Phi_{\mathbb{Q}}^{D_4}(4) = \Phi_{\mathbb{Q}}(2) \cup \{ \mathcal{C}_{20}, \mathcal{C}_{24} \}$.

This will complete the determination of $\Phi_{\mathbb{Q}}(4)$.

Corollary 8.7. *The set $\Phi_{\mathbb{Q}}(4)$ is given by*

$$\Phi_{\mathbb{Q}}(4) = \{C_n : n = 1, \dots, 10, 12, 13, 15, 16, 20, 24\} \cup \{C_2 \times C_{2m} : m = 1, \dots, 6, 8\} \\ \cup \{C_3 \times C_{3m} : m = 1, 2\} \cup \{C_4 \times C_{4m} : m = 1, 2\} \cup \{C_5 \times C_5\} \cup \{C_6 \times C_6\}.$$

Remark 8.8. Corollary 8.7 implies that the list of possible pairs $(E(\mathbb{Q})_{\text{tors}}, E(K)_{\text{tors}})$, where K is a quartic field, obtained by the first author and Lozano-Robledo in [13] is complete.

Throughout this section, denote by K a quartic number field whose normal closure \widehat{K} over \mathbb{Q} has a Galois group isomorphic to D_4 . We will say that such a quartic number field is a *dihedral quartic number field*. Denote by F the unique quadratic field contained in K . Elliptic curves will always, unless stated otherwise, be defined over \mathbb{Q} .

We will make use of the following lemma.

Lemma 8.9. *Let p be a prime, let E/F be an elliptic curve over a number field F , and let L/F be a finite extension such that $\text{Gal}(\widehat{L}/F)$ does not contain as quotient a cyclic group of order $a = [F(\zeta_p) : F]$, where \widehat{L} is the normal closure of L over F .*

Then if $E(L)$ has a point P of order p , it follows that E has a p -isogeny over F . Furthermore $F(P)$ is Galois over F and $\text{Gal}(F(P)/F)$ is a cyclic group of order dividing $p - 1$.

Proof. Suppose $E(L)$ contains a point P of order p . By our assumptions, \widehat{L} does not contain $\mathbb{Q}(\zeta_p)$, since $\text{Gal}(F(\zeta_p)/F) \simeq C_a$ would be a quotient of $\text{Gal}(\widehat{L}/F)$, contradicting our assumptions. Hence $E(\widehat{L}) \simeq C_p$, and since \widehat{L} is Galois over F , it follows that $\langle P \rangle$ is a cyclic Galois-invariant subgroup, and hence E has a p -isogeny ϕ over F such that $\langle P \rangle = \ker \phi$. Since P is in the kernel of an F -rational isogeny, it follows that $F(P)$ is Galois over F of degree dividing $p - 1$, over F . \square

8.1.1. *p -primary torsion subgroup.* The first step in determining $\Phi_{\mathbb{Q}}^{D_4}(4)$ is to determine the possibilities for $E(K)[p^\infty]$ for all primes p .

Lemma 8.10. *If P is a point of prime order p in $E(K)$, then $p \in \{2, 3, 5, 7\}$.*

Proof. First note that $R_{\mathbb{Q}}(2) = \{2, 3, 5, 7\}$, therefore the primes in the above set appear as orders of points over dihedral quartic number fields. By Corollary 6.1, we have $R_{\mathbb{Q}}(4) = \{2, 3, 5, 7, 13\}$.

We claim that there are no points of order 13 over dihedral quartic number fields. Suppose the opposite; let $P \in E(K)$ be a point of order 13. By Lemma 8.9, $\mathbb{Q}(P)$ is Galois over \mathbb{Q} , and hence $\mathbb{Q}(P) = \mathbb{Q}$ or $\mathbb{Q}(P) = F$. But that is in contradiction with the fact that $R_{\mathbb{Q}}(2) = \{2, 3, 5, 7\}$. \square

Lemma 8.11. *The order of $E(K)[p^\infty]$ divides 16, 9, 5, 7, for $p = 2, 3, 5, 7$, respectively. In particular all the possible orders occur.*

Proof. Let n be a positive integer such that $n_p \leq 4, 2, 1, 1$, if $p = 2, 3, 5, 7$, respectively, then $C_{p^{n_p}} \in \Phi(1) \subseteq \Phi_{\mathbb{Q}}^{D_4}(4)$.

We split the proof depending on the prime p :

- $p = 2$: The possible torsion subgroups of elliptic curve with CM defined over a quartic number field have been determined in [8]. In particular, it is proven that the order of the 2-primary torsion subgroup divides 16. Therefore we can assume E is non-CM. The minimal degree of definition of any 2-subgroup of $E(\overline{\mathbb{Q}})$ has been

determined in [12, Theorem 1.4] in the non-CM case. In particular, for degree 4 we have (see Table 1 in [12]) that if E/\mathbb{Q} is a non-CM elliptic curve and K/\mathbb{Q} is a quartic number field, then

$$E(K)[2^\infty] \in \{ \mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_4, \mathcal{C}_8, \mathcal{C}_{16}, \mathcal{C}_2 \times \mathcal{C}_2, \mathcal{C}_2 \times \mathcal{C}_4, \mathcal{C}_2 \times \mathcal{C}_8, \\ \mathcal{C}_2 \times \mathcal{C}_{16}, \mathcal{C}_4 \times \mathcal{C}_4, \mathcal{C}_4 \times \mathcal{C}_8 \}.$$

We will prove that the groups $\mathcal{C}_2 \times \mathcal{C}_{16}$ and $\mathcal{C}_4 \times \mathcal{C}_8$ do not occur over dihedral quartic number fields.

Rouse and Zureick-Brown [33] have classified all the possible 2-adic images of $\rho_{E,2^\infty} : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow \text{GL}_2(\mathbb{Z}_2)$, and have given explicitly all the 1208 possibilities. The first author and Lozano-Robledo [12] have determined for each possible image the degree of the field of definition of any 2-subgroup. From the file `2primary_Ss.txt` (this file can be found at the research website of the first author [12]) one can read out if a given 2-subgroup is defined over a number field of given degree d . In particular, the 2-adic images up to conjugation (following the notation of [33]) for the case $d = 4$ and the given 2-subgroups are:

- $\mathcal{C}_2 \times \mathcal{C}_{16}$: X193j, X193l, X193n, X213i, X213k, X215c, X215d, X235f, X235l, X235m,
- $\mathcal{C}_4 \times \mathcal{C}_8$: X183d, X183g, X183i, X187d, X187h, X187j, X187k, X189b, X189d, X189e, X193g, X193i, X193n, X194g, X194h, X194k, X194l, X195h, X195j, X195l.

For each of these images, we determine (see [14]) the Galois group of the quartic number field over \mathbb{Q} corresponding to each of the above 2-adic images (i.e., the possible Galois groups are the normal cores of index 4 stabilizers of the appropriate element in $(\mathbb{Z}/2^n\mathbb{Z})^2$). We obtain that all the quartic number fields are Galois over \mathbb{Q} . This finishes the proof for the 2-primary torsion.

• $p = 3$: We have $\mathcal{C}_3 \times \mathcal{C}_3 \in \Phi_{\mathbb{Q}}(2)$, therefore it occurs as the torsion subgroup of elliptic curves over dihedral quartic number fields. By [13, Theorem 7 (11)] we have that $\mathcal{C}_3 \times \mathcal{C}_9$ is not a subgroup of $E(K)$. Thus, to finish determining the possible 3-primary torsion it is enough to prove that there is no 27-torsion over a dihedral quartic number field K . Suppose the opposite, i.e., \mathcal{C}_{27} is a subgroup of $E(K)$. Since F is a subfield of K and $[K : F] = 2$, there exists a $\delta \in F$ such that $K = F(\sqrt{\delta})$. Then we have

$$E(K)[27] \simeq E(F)[27] \oplus E^\delta(F)[27].$$

Therefore $E(F)[27]$ or $E^\delta(F)[27]$ contains \mathcal{C}_{27} . But this is not possible since \mathcal{C}_{27} is not a subgroup of any group in $\Phi(2)$.

• $p = 5$: It is not possible to have the full 5-torsion over a dihedral quartic number field K , since otherwise $\mathbb{Q}(\zeta_5) = K$, and this is impossible since $\mathbb{Q}(\zeta_5)$ is cyclic. We observe that \mathcal{C}_{25} is not possible over dihedral quartic number fields by using the same argument as in the \mathcal{C}_{27} case.

• $p = 7$: It is impossible that $E[7] \subseteq E(K)$ since $K \not\supseteq \mathbb{Q}(\zeta_7)$. To prove that there is no 49-torsion, we again use the same argument as in the \mathcal{C}_{27} case. □

8.1.2. *Combinations of p -primary torsion.* In the previous section we have determined the p -primary torsion of an elliptic curve E/\mathbb{Q} over a dihedral quartic number

field. In this section we study the possible combinations of p -power torsion and q -power torsion for primes $p \neq q$. First we prove that if $E(K)$ has a point of order 7, then $E(K)[p] = \{\mathcal{O}\}$ for all primes $p \neq 7$.

Lemma 8.12. *If 7 divides the order of $E(K)_{\text{tors}}$, then $E(K)_{\text{tors}} \simeq \mathcal{C}_7$.*

Proof. Suppose $E(K)_{\text{tors}}$ has a point of order 7. Let p be a prime, $p \neq 7$ such that p divides the order of $E(K)_{\text{tors}}$. By Lemma 8.10 we have that $p \in \{2, 3, 5\}$. The case $p = 2$ is not possible, since there cannot be any 14-torsion over a quartic number field (the proof of [7, Proposition 3.9] can be slightly modified to prove it). The case $p = 3$ is also not possible, since there cannot exist points of order 21 over a quartic number field (see [13, Theorem 7 (iv)]). Finally, suppose $p = 5$. Then we have $E(\widehat{K})[35] \simeq \mathcal{C}_{35}$, because \widehat{K} does not contain $\mathbb{Q}(\zeta_5)$ or $\mathbb{Q}(\zeta_7)$, as \mathcal{C}_4 and \mathcal{C}_6 are not quotients of D_4 . This implies that E has a 35-isogeny over \mathbb{Q} which is not possible by Theorem 2.1.

Now, by Lemma 8.11, we deduce that $E(K)_{\text{tors}} \simeq \mathcal{C}_7$. □

Lemma 8.13. *If 15 divides the order of $E(K)_{\text{tors}}$, then $E(K)_{\text{tors}} \simeq \mathcal{C}_{15}$.*

Proof. We have that $\mathcal{C}_{15} \in \Phi_{\mathbb{Q}}(2)$, therefore the case $E(K)_{\text{tors}} \simeq \mathcal{C}_{15}$ is possible. Now, let $P \in E(K)_{\text{tors}}$ be a point of order 15. Theorem 8 in [13] shows that $\mathbb{Q}(P)$ is an abelian extension. Thus, since K is a dihedral quartic number field, we have that $\mathbb{Q}(P) = F$, where F is the unique quadratic subfield of K .

If there existed a point R of order 30 in $E(K)_{\text{tors}}$, then $2R$ would be a point of order 15 and $\mathbb{Q}(2R) = F$ as above. The point R is not defined over F , as $\mathcal{C}_{15} \notin \Phi_{\mathbb{Q}}(2)$. But then $[\mathbb{Q}(R) : \mathbb{Q}(2R)] = 3$, which is a contradiction. We conclude that $E(K)_{\text{tors}}$ cannot contain a point of order 30.

Now by Lemma 8.11 we have that the 3-primary torsion is isomorphic to one of the groups: $\mathcal{C}_3, \mathcal{C}_9, \mathcal{C}_3 \times \mathcal{C}_3$. To finish the proof we prove that 9 does not divide the order of $E(K)_{\text{tors}}$. If the 3-primary torsion is isomorphic to \mathcal{C}_9 , we have that any point of order 9 is defined over a proper subfield of K (see Theorem 7 (10) from [13]). By the previous argument we deduce that E has a point of order 45 over the quadratic field F , which is not possible. Finally, let us suppose that the $\mathcal{C}_{15} \times \mathcal{C}_3$ is a subgroup of $E(K)_{\text{tors}}$. Then the quadratic subfield of K is $\mathbb{Q}(\sqrt{-3})$, and we claim that both the subgroups isomorphic to \mathcal{C}_5 and to $\mathcal{C}_3 \times \mathcal{C}_3$ have to be defined over $\mathbb{Q}(\sqrt{-3})$, which is clearly impossible. For the subgroup \mathcal{C}_5 , this follows from Lemma 8.9, while for $\mathcal{C}_3 \times \mathcal{C}_3$ it follows from the fact that $\mathbb{Q}(E[3])$ is Galois over \mathbb{Q} . □

Lemma 8.14. *If 10 divides the order of $E(K)_{\text{tors}}$, then $E(K)_{\text{tors}}$ is isomorphic to either $\mathcal{C}_{10}, \mathcal{C}_{20}$ or to $\mathcal{C}_2 \times \mathcal{C}_{10}$.*

Proof. Let us prove that $\mathcal{C}_{40}, \mathcal{C}_2 \times \mathcal{C}_{20} \subsetneq E(K)_{\text{tors}}$. Suppose first that $E(K)_{\text{tors}} \supseteq \mathcal{C}_2 \times \mathcal{C}_{20}$. By Lemma 8.9, we have that E has a 5-isogeny over \mathbb{Q} and that $\mathbb{Q}(P)$ is Galois over \mathbb{Q} . Thus it follows that $\mathbb{Q}(P) = F$ or $\mathbb{Q}(P) = \mathbb{Q}$. If $\mathbb{Q}(P) = F = \mathbb{Q}(\sqrt{d})$, then we have $E(F)[5] = E(\mathbb{Q})[5] \oplus E^d(\mathbb{Q})[5]$. Then either $E(\mathbb{Q})[5] \simeq \mathcal{C}_5$ or $E^d(\mathbb{Q})[5] \simeq \mathcal{C}_5$. In the latter case we can study E^d instead of E ; thus it follows that we can take without loss of generality that $\mathbb{Q}(P) = \mathbb{Q}$. We conclude that $E(\mathbb{Q})_{\text{tors}} \simeq \mathcal{C}_{10}$. Let $Q \in E(\mathbb{Q})$ be a point of order 10. The torsion of E will grow from \mathcal{C}_{10} in \mathbb{Q} to $\mathcal{C}_2 \times \mathcal{C}_{20}$, if and only if a solution R of the equation $2R = Q$ satisfies that $\mathbb{Q}(R)$ contains the quadratic field $\mathbb{Q}(\sqrt{\Delta_E}) = \mathbb{Q}(E[2])$ as a subfield.

The universal elliptic curve over $X_1(10)$ (which can be thought of as the parametric family of elliptic curves with \mathcal{C}_{10} torsion) is (see [26]):

$$E_t : y^2 + (t^3 - 2t^2 - 4t + 4)xy + (t - 1)(t - 2)(t^2 - 6t + 4)t^3y = x^3 - (t - 1)(t - 2)t^3x^2,$$

where we can take $Q = (0, 0) \in E_t[10]$. Note that the four solutions of $2R = Q$ are defined over at most 2 number fields, but they will be conjugate to each other, and hence the following argumentation does not depend on the choice of R . The condition that $\mathbb{Q}(\sqrt{\Delta_E})$ is contained in $\mathbb{Q}(R)$ is, after some calculation, equivalent to the hyperelliptic curve

$$C : y^2 = x(x - 1)(x - 2)(x^2 - x - 1)(x^2 - 6x + 4)$$

having a solution such that $y \neq 0$. To find all the rational points on C , we note that $\text{Aut}(C) \simeq D_4$ and for a non-hyperelliptic involution $\sigma \in \text{Aut}(C)$ we obtain (see [14]) that the quotient curve $X = C/\langle\sigma\rangle$ is of genus 1, where X is defined by the following equation:

$$X : y^2 = x^3 - 7x - 6.$$

We have $X(\mathbb{Q}) = \{\mathcal{O}, (-1, 0), (-2, 0), (3, 0)\} \simeq \mathcal{C}_2 \times \mathcal{C}_2$ and the map $\phi : C \rightarrow X$ sends the Weierstrass points of C to the 2-torsion points of X . We obtain (see [14]) that the 4 rational Weierstrass points on C , $\{\infty, (0, 0), (1, 0), (2, 0)\}$ map to $\{\mathcal{O}, (-1, 0)\}$; the preimages of the 2 remaining points in $X(\mathbb{Q})$ are the 4 quadratic Weierstrass points of C . This proves that

$$C(\mathbb{Q}) = \{\infty, (0, 0), (1, 0), (2, 0)\}.$$

Thus we have proved that $E(K)$ cannot contain $\mathcal{C}_2 \times \mathcal{C}_{20}$.

Suppose now that $E(K)_{\text{tors}} \supseteq \mathcal{C}_{40}$. By the same argument as above, we can suppose that $E(\mathbb{Q})$ contains a point of order 5 and that $E(\mathbb{Q})_{\text{tors}} \simeq \mathcal{C}_{10}$. Also, the 2-Sylow subgroup of the torsion is \mathcal{C}_2 over \mathbb{Q} , \mathcal{C}_8 over K , and \mathcal{C}_2 over the quadratic subfield of K (as otherwise E would have a 20-torsion point over a quadratic field). From the file `2primary_Ss.txt` [12] we see (see [14]) that the 2-adic Galois representation of E has to be **X191**, using the notation from [33]. The modular curve **X191** covers the modular curve **X11**, so our elliptic curve has to correspond to a point on $\mathbf{X11} \times_{X_0(1)} X_0(5)$. By studying the j -maps of **X11** and $X_0(5)$, we get an elliptic curve C (with Cremona label **80a1**) with rank 0 over \mathbb{Q} and 4 rational points. We check (see [14]) that none of the points correspond to an elliptic curve, completing the proof. \square

Lemma 8.15. *If 6 divides the order of $E(K)_{\text{tors}}$, then $E(K)_{\text{tors}}$ is isomorphic to either \mathcal{C}_6 , \mathcal{C}_{12} , \mathcal{C}_{24} , $\mathcal{C}_2 \times \mathcal{C}_6$, $\mathcal{C}_2 \times \mathcal{C}_{12}$, or $\mathcal{C}_3 \times \mathcal{C}_6$.*

Proof. By [13, Theorem 7 (11)], we see that $E(K)_{\text{tors}}$ cannot contain \mathcal{C}_{18} .

We can see that $E(K)_{\text{tors}}$ does not contain $\mathcal{C}_6 \times \mathcal{C}_6$, as $\mathbb{Q}(E[6])$ is a Galois extension of \mathbb{Q} , hence it would have to be either \mathbb{Q} or F . But it is impossible to have full 6-torsion over \mathbb{Q} or over a quadratic field.

By [1, Theorem 8], we see that $\mathcal{C}_3 \times \mathcal{C}_{12}$ and $\mathcal{C}_4 \times \mathcal{C}_{12}$ cannot be contained in $E(K)_{\text{tors}}$.

We claim that $E(K)_{\text{tors}} \supseteq \mathcal{C}_2 \times \mathcal{C}_{24}$ is impossible. By a search (see [14]) of the database of 2-adic images [33], we obtain that the 2-power torsion of E cannot grow from \mathcal{C}_2 over \mathbb{Q} to $\mathcal{C}_2 \times \mathcal{C}_8$ in a D_4 extension.

It follows that $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_2$ or $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_4$.

Now we split the proof in two cases: either $G_E(3)$ is conjugate to 3Ns or not.

We will show that if $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_2$ or $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_4$, then $G_E(3)$ is not conjugate to 3Ns. By [37, Theorem 1.1], $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_2$ if and only if

$$j(E) = 256 \frac{(t^2 + t + 1)^3}{t^2(t+1)^2} \quad \text{for some } t \in \mathbb{Q} \setminus \{0, -1\}.$$

On the other hand, if $G_E(3)$ is conjugate in $\text{GL}_2(\mathbb{F}_3)$ to 3Ns, then by [37, Theorem 1.2], we have

$$j(E) = s^3 \quad \text{for some } s \in \mathbb{Q}.$$

Hence we obtain the equation

$$C : s^3 = 256 \frac{(t^2 + t + 1)^3}{t^2(t+1)^2},$$

which cuts out a singular genus 1 curve, which is birationally equivalent to

$$X_1 : y^2 = x^3 + 1,$$

which has rank 0 and 6 rational points. We obtain (see [14]) that the only (affine) rational points on X_1 satisfy $s = 12$, implying $j(E) = 1728$. But by [8, §4.4], elliptic curves with CM cannot have $\mathcal{C}_2 \times \mathcal{C}_{24}$ torsion over quartic number fields.

If $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_4$, then E has a 4-isogeny over \mathbb{Q} , from which it follows that [28, Table 3]

$$j(E) = \frac{(h^2 + 16h + 16)^3}{h(h+16)} \quad \text{for some } h \in \mathbb{Q} \setminus \{0, -16\}.$$

Thus, $h(h+16)$ needs to be a cube, giving an equation of an elliptic curve

$$X_2 : h^2 + 16h = v^3.$$

The elliptic curve X_2 has (see [14]) rank 0 over \mathbb{Q} and its torsion points (with the assumption $h \neq 0, -16$) give $j(E) = 1728$ and 287496, both of which are CM values. Again, by [8, §4.4], we note that elliptic curves with CM cannot have $\mathcal{C}_2 \times \mathcal{C}_{24}$ torsion over quartic number fields, which rules out this possibility.

Thus, we have proved $G_E(3)$ is not conjugate to 3Ns. We claim that then $E(F)[3] \neq \{\mathcal{O}\}$. To see this, in Table 1, we see that the only possibility of $\mathbb{Q}(P)$ being equal to K (which is equivalent to $d_v = 4$ for some v) is that $G_E(3)$ is conjugate to 3Cs. But if $G_E(3)$ is 3Cs and $[\mathbb{Q}(P) : \mathbb{Q}] = 4$ for a point P of order 3, then it would follow, since $\mathbb{Q}(P) \subseteq \mathbb{Q}(E[3])$ and both $\mathbb{Q}(E[3])$ and $\mathbb{Q}(P)$ are of degree 4, that $\mathbb{Q}(E[3]) = \mathbb{Q}(P) = K$. But $\mathbb{Q}(E[3])$ is Galois over \mathbb{Q} and K is not. We conclude that $E(F)[3] \neq \{\mathcal{O}\}$.

We now claim that if $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_2$, then $E(K) \supseteq \mathcal{C}_2 \times \mathcal{C}_{24}$ is impossible. Since $E(F)[3] = E(\mathbb{Q})[3] \oplus E^d(\mathbb{Q})[3]$, where $F = \mathbb{Q}(\sqrt{d})$, so either E or E^d has a 3-torsion point; we choose without loss of generality E to be the twist with 3-torsion. Since quadratic twisting does not change the 2-torsion, we have that $E(\mathbb{Q}) \supseteq \mathcal{C}_2 \times \mathcal{C}_6$. By a search (see [14]) of the database of 2-adic images [33], we find that there exist

2-adic representations such that 2-power torsion of E grows from $\mathcal{C}_2 \times \mathcal{C}_2$ over \mathbb{Q} to $\mathcal{C}_2 \times \mathcal{C}_8$ over F or/and over K . In the former case we obtain $\mathcal{C}_2 \times \mathcal{C}_{24} \subseteq E(F)$, which is impossible over quadratic fields. In the last case, $\mathcal{C}_2 \times \mathcal{C}_8 \subset E(K)[2^\infty]$, we obtain that the field $L = \mathbb{Q}(E[4])$ is a quartic number field. But together with our assumption that $E(\mathbb{Q})[3] \neq 0$, we have that $E(L)$ contains a subgroup isomorphic to $\mathcal{C}_4 \times \mathcal{C}_{12}$, which is in contradiction with [1, Theorem 7].

It remains to prove that if $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_4$, then $E(K)$ cannot contain $\mathcal{C}_2 \times \mathcal{C}_{24}$. By a search (see [14]) of the database of 2-adic images [33], we find that if $E(\mathbb{Q})[2^\infty] \supseteq \mathcal{C}_4$ and $E(K)[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_8$, then it is necessary that already $E(F)[2^\infty] \supseteq \mathcal{C}_2 \times \mathcal{C}_8$. But this implies that $E(F) \supseteq \mathcal{C}_2 \times \mathcal{C}_{24}$, which is impossible over quadratic number fields.

Thus we have proved that $\mathcal{C}_2 \times \mathcal{C}_{24}$ torsion is impossible over K .

Suppose now $E(K) \supseteq \mathcal{C}_{48}$. By a search in the file `2primary_Ss.txt` [12] we find (see [14]) that if $E(K)[2^\infty] \supseteq \mathcal{C}_{16}$, then $E(F)[2^\infty] \supseteq \mathcal{C}_8$, where F is the quadratic subfield of K . This implies that E has an 8-isogeny over \mathbb{Q} . Since E cannot have both a 3-isogeny and an 8-isogeny, using the same argumentation as in the previous case, it follows that $G_E(3)$ has to be $3Ns$. But we have already shown that E cannot have both a 4-isogeny and $G_E(3)$ to be $3Ns$, completing the proof. \square

Proof of Theorem 8.5. Finally, to prove Theorem 8.5, we note that all the groups that are in $\Phi_{\mathbb{Q}}(2)$ are also in $\Phi_{\mathbb{Q}}^{D_4}(4)$ and by [19], there exists infinitely many elliptic curves with \mathcal{C}_{20} and \mathcal{C}_{24} torsion over dihedral quartic number fields. Therefore $\Phi_{\mathbb{Q}}(2) \cup \{\mathcal{C}_{20}, \mathcal{C}_{24}\} \subseteq \Phi_{\mathbb{Q}}^{D_4}(4)$. The equality holds by Lemmas 8.11-8.15. \square

Proof of Corollary 8.7. Let K a quartic number field; then the group $\text{Gal}(\widehat{K}/\mathbb{Q})$ is isomorphic to S_4, A_4, D_4, V_4 , or \mathcal{C}_4 . Therefore

$$\Phi_{\mathbb{Q}}(4) = \Phi_{\mathbb{Q}}^{S_4}(4) \cup \Phi_{\mathbb{Q}}^{A_4}(4) \cup \Phi_{\mathbb{Q}}^{D_4}(4) \cup \Phi_{\mathbb{Q}}^{V_4}(4) \cup \Phi_{\mathbb{Q}}^{\mathcal{C}_4}(4).$$

The cases $\Phi_{\mathbb{Q}}^{V_4}(4)$ and $\Phi_{\mathbb{Q}}^{\mathcal{C}_4}(4)$ have been determined by Chou [7]. The remaining cases are dealt with in Corollaries 8.4 and 8.7. \square

APPENDIX: IMAGES OF MOD p GALOIS REPRESENTATIONS
ASSOCIATED TO ELLIPTIC CURVES OVER \mathbb{Q}

For each possible known subgroup $G_E(p) \subsetneq \text{GL}_2(\mathbb{F}_p)$ where E/\mathbb{Q} is a non-CM elliptic curve and p is a prime, Tables 1 and 2 list in the first and second columns the corresponding labels in Sutherland and Zywina notation, and the following data:

- $d_v = [G_E(p) : G_E(p)_v] = |G_E(p).v|$ for $v \in \mathbb{F}_p^2, v \neq (0, 0)$; equivalently, the degrees of the extensions $\mathbb{Q}(P)$ over \mathbb{Q} for points $P \in E(\overline{\mathbb{Q}})$ of order p .
- $d = |G_E(p)|$; equivalently, the degree $\mathbb{Q}(E[p])$ over \mathbb{Q} .

Note that Tables 1 and 2 are partially extracted from Table 3 of [36]. The difference is that [36, Table 3] only lists the minimum of d_v , which is denoted by d_1 therein. The computations (see [14]) needed to obtain these tables were performed in Magma.

TABLE 1. Possible images $G_E(p) \neq \text{GL}_2(\mathbb{F}_p)$, for $p \leq 11$, for non-CM elliptic curves E/\mathbb{Q} .

| Sutherland | Zywina | d_v | d | Sutherland | Zywina | d_v | d |
|------------|-----------|---------|-----|------------|-----------|----------|-----|
| 2Cs | G_1 | 1 | 1 | 7Ns. 2.1 | $H_{1,1}$ | 6, 9, 18 | 18 |
| 2B | G_2 | 1, 2 | 2 | 7Ns. 3.1 | G_1 | 12, 18 | 36 |
| 2Cn | G_3 | 3 | 3 | 7B. 1.1 | $H_{3,1}$ | 1, 42 | 42 |
| 3Cs. 1.1 | $H_{1,1}$ | 1, 2 | 2 | 7B. 1.3 | $H_{4,1}$ | 6, 7 | 42 |
| 3Cs | G_1 | 2, 4 | 4 | 7B. 1.2 | $H_{5,2}$ | 3, 42 | 42 |
| 3B. 1.1 | $H_{3,1}$ | 1, 6 | 6 | 7B. 1.5 | $H_{5,1}$ | 6, 21 | 42 |
| 3B. 1.2 | $H_{3,2}$ | 2, 3 | 6 | 7B. 1.6 | $H_{3,2}$ | 2, 21 | 42 |
| 3Ns | G_2 | 4 | 8 | 7B. 1.4 | $H_{4,2}$ | 3, 14 | 42 |
| 3B | G_3 | 2, 6 | 12 | 7Ns | G_2 | 12, 36 | 72 |
| 3Nn | G_4 | 8 | 16 | 7B. 6.1 | G_3 | 2, 42 | 84 |
| 5Cs. 1.1 | $H_{1,1}$ | 1, 4 | 4 | 7B. 6.3 | G_4 | 6, 14 | 84 |
| 5Cs. 1.3 | $H_{1,2}$ | 2, 4 | 4 | 7B. 6.2 | G_5 | 6, 42 | 84 |
| 5Cs. 4.1 | G_1 | 2, 4, 8 | 8 | 7Nn | G_6 | 48 | 96 |
| 5Ns. 2.1 | G_3 | 8, 16 | 16 | 7B. 2.1 | $H_{7,2}$ | 3, 42 | 126 |
| 5Cs | G_2 | 4, 4 | 16 | 7B. 2.3 | $H_{7,1}$ | 6, 21 | 126 |
| 5B. 1.1 | $H_{6,1}$ | 1, 20 | 20 | 7B | G_7 | 6, 42 | 252 |
| 5B. 1.2 | $H_{5,1}$ | 4, 5 | 20 | 11B. 1.4 | $H_{1,1}$ | 5, 110 | 110 |
| 5B. 1.4 | $H_{6,2}$ | 2, 20 | 20 | 11B. 1.5 | $H_{2,1}$ | 5, 110 | 110 |
| 5B. 1.3 | $H_{5,2}$ | 4, 10 | 20 | 11B. 1.6 | $H_{2,2}$ | 10, 55 | 110 |
| 5Ns | G_4 | 8, 16 | 32 | 11B. 1.7 | $H_{1,2}$ | 10, 55 | 110 |
| 5B. 4.1 | G_6 | 2, 20 | 40 | 11B. 10.4 | G_1 | 10, 110 | 220 |
| 5B. 4.2 | G_5 | 4, 10 | 40 | 11B. 10.5 | G_2 | 10, 110 | 220 |
| 5Nn | G_7 | 24 | 48 | 11Nn | G_3 | 120 | 240 |
| 5B | G_8 | 4, 20 | 80 | | | | |
| 5S4 | G_9 | 24 | 96 | | | | |

TABLE 2. Known images $G_E(p) \neq \text{GL}_2(\mathbb{F}_p)$, for $p = 13, 17$, or 37 , for non-CM elliptic curves E/\mathbb{Q} .

| Sutherland | Zywina | d_v | d | Sutherland | Zywina | d_v | d |
|------------|-----------|---------|------|------------|--------|----------|-------|
| 13S4 | G_7 | 72, 96 | 288 | 17B. 4.2 | G_1 | 8, 272 | 1088 |
| 13B. 3.1 | $H_{5,1}$ | 3, 156 | 468 | 17B. 4.6 | G_2 | 16, 136 | 1088 |
| 13B. 3.2 | $H_{4,1}$ | 12, 39 | 468 | 37B. 8.1 | G_1 | 12, 1332 | 15984 |
| 13B. 3.4 | $H_{5,2}$ | 6, 156 | 468 | 37B. 8.2 | G_2 | 36, 444 | 15984 |
| 13B. 3.7 | $H_{4,2}$ | 12, 78 | 468 | | | | |
| 13B. 5.1 | G_2 | 4, 156 | 624 | | | | |
| 13B. 5.2 | G_1 | 12, 52 | 624 | | | | |
| 13B. 5.4 | G_3 | 12, 156 | 624 | | | | |
| 13B. 4.1 | G_5 | 6, 156 | 936 | | | | |
| 13B. 4.2 | G_4 | 12, 78 | 936 | | | | |
| 13B | G_6 | 12, 156 | 1872 | | | | |

ACKNOWLEDGMENTS

We thank Abbey Bourdon for suggesting Remark 7.5, Pete Clark and Tomislav Gužvić for pointing out some mistakes in a previous version of the paper, Maarten Derickx for pointing out the example in Remark 4.7, and the anonymous referee for helpful comments.

REFERENCES

- [1] P. Bruin and F. Najman, *A criterion to rule out torsion groups for elliptic curves over number fields*, Res. Number Theory **2** (2016), Art. 3, 13, DOI 10.1007/s40993-015-0031-5. MR3501016 ↑1480, 1482
- [2] J. Balakrishnan, N. Dogra, J. S. Müller, J. Tuitman, and J. Vonk, *Explicit Chabauty–Kim for the split Cartan modular curve of level 13*, Ann. of Math. (2) **189** (2019), no. 3, 885–944, DOI 10.4007/annals.2019.189.3.6. MR3961086 ↑1461
- [3] Y. Bilu and P. Parent, *Serre’s uniformity problem in the split Cartan case*, Ann. of Math. (2) **173** (2011), no. 1, 569–584, DOI 10.4007/annals.2011.173.1.13. MR2753610 ↑1461
- [4] Y. Bilu, P. Parent, and M. Rebolledo, *Rational points on $X_0^+(p^r)$* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **63** (2013), no. 3, 957–984, DOI 10.5802/aif.2781. MR3137477 ↑1461
- [5] W. Bosma, J. J. Cannon, C. Fieker, and A. Steel (eds.), *Handbook of Magma Functions*, Edition 2.21 (2015). 1459
- [6] A. Bourdon, P. L. Clark, and J. Stankewicz, *Torsion points on CM elliptic curves over real number fields*, Trans. Amer. Math. Soc. **369** (2017), no. 12, 8457–8496, DOI 10.1090/tran/6905. MR3710632 ↑1472
- [7] M. Chou, *Torsion of rational elliptic curves over quartic Galois number fields*, J. Number Theory **160** (2016), 603–628, DOI 10.1016/j.jnt.2015.09.013. MR3425225 ↑1475, 1479, 1482
- [8] P. L. Clark, P. Corn, A. Rice, and J. Stankewicz, *Computation on elliptic curves with complex multiplication*, LMS J. Comput. Math. **17** (2014), no. 1, 509–535, DOI 10.1112/S1461157014000072. MR3356044 ↑1472, 1474, 1477, 1481
- [9] M. Derickx and M. van Hoeij, *Gonality of the modular curve $X_1(N)$* , J. Algebra **417** (2014), 52–71, DOI 10.1016/j.jalgebra.2014.06.026. MR3244637 ↑1459
- [10] M. Derickx and A. V. Sutherland, *Torsion subgroups of elliptic curves over quintic and sextic number fields*, Proc. Amer. Math. Soc. **145** (2017), no. 10, 4233–4245, DOI 10.1090/proc/13605. MR3690609 ↑1458
- [11] E. González-Jiménez, *Complete classification of the torsion structures of rational elliptic curves over quintic number fields*, J. Algebra **478** (2017), 484–505, DOI 10.1016/j.jalgebra.2017.01.012. MR3621686 ↑1458, 1474, 1475
- [12] E. González-Jiménez and Á. Lozano-Robledo, *On the minimal degree of definition of p -primary torsion subgroups of elliptic curves*, Math. Res. Lett. **24** (2017), no. 4, 1067–1096, DOI 10.4310/MRL.2017.v24.n4.a7. MR3723804 ↑1478, 1480, 1482
- [13] E. González-Jiménez and Á. Lozano-Robledo, *On the torsion of rational elliptic curves over quartic fields*, Math. Comp. **87** (2018), no. 311, 1457–1478, DOI 10.1090/mcom/3235. MR3766394 ↑1477, 1478, 1479, 1480
- [14] E. González-Jiménez and F. Najman, *Magma scripts and electronic transcript of computations for the paper, Growth of torsion groups of elliptic curves upon base change*, <http://matematicas.uam.es/~enrique.gonzalez.jimenez> 1459, 1467, 1468, 1470, 1471, 1472, 1473, 1475, 1476, 1478, 1480, 1481, 1482
- [15] E. González-Jiménez, F. Najman, and J. M. Tornero, *Torsion of rational elliptic curves over cubic fields*, Rocky Mountain J. Math. **46** (2016), no. 6, 1899–1917, DOI 10.1216/RMJ-2016-46-6-1899. MR3591265 ↑1474
- [16] E. González-Jiménez and J. M. Tornero, *Torsion of rational elliptic curves over quadratic fields II*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **110** (2016), no. 1, 121–143, DOI 10.1007/s13398-015-0223-9. MR3462078 ↑1474
- [17] D. Jeon, C. H. Kim, and A. Schweizer, *On the torsion of elliptic curves over cubic number fields*, Acta Arith. **113** (2004), no. 3, 291–301, DOI 10.4064/aa113-3-6. MR2069117 ↑1458
- [18] D. Jeon, C. H. Kim, and E. Park, *On the torsion of elliptic curves over quartic number fields*, J. London Math. Soc. (2) **74** (2006), no. 1, 1–12, DOI 10.1112/S0024610706022940. MR2254548 ↑1458
- [19] D. Jeon, C. H. Kim, and Y. Lee, *Infinite families of elliptic curves over dihedral quartic number fields*, J. Number Theory **133** (2013), no. 1, 115–122, DOI 10.1016/j.jnt.2012.06.014. MR2981403 ↑1482
- [20] S. Kamienny, *Torsion points on elliptic curves and q -coefficients of modular forms*, Invent. Math. **109** (1992), no. 2, 221–229, DOI 10.1007/BF01232025. MR1172689 ↑1458, 1460

- [21] M. A. Kenku, *The modular curve $X_0(39)$ and rational isogeny*, Math. Proc. Cambridge Philos. Soc. **85** (1979), no. 1, 21–23, DOI 10.1017/S0305004100055444. MR510395 ↑1459
- [22] M. A. Kenku, *The modular curves $X_0(65)$ and $X_0(91)$ and rational isogeny*, Math. Proc. Cambridge Philos. Soc. **87** (1980), no. 1, 15–20, DOI 10.1017/S0305004100056462. MR549292 ↑1459
- [23] M. A. Kenku, *The modular curve $X_0(169)$ and rational isogeny*, J. London Math. Soc. (2) **22** (1980), no. 2, 239–244, DOI 10.1112/jlms/s2-22.2.239. MR588271 ↑1459
- [24] M. A. Kenku, *On the modular curves $X_0(125)$, $X_1(25)$ and $X_1(49)$* , J. London Math. Soc. (2) **23** (1981), no. 3, 415–427, DOI 10.1112/jlms/s2-23.3.415. MR616546 ↑1459
- [25] M. A. Kenku and F. Momose, *Torsion points on elliptic curves defined over quadratic fields*, Nagoya Math. J. **109** (1988), 125–149, DOI 10.1017/S0027763000002816. MR931956 ↑1458, 1460
- [26] D. S. Kubert, *Universal bounds on the torsion of elliptic curves*, Proc. London Math. Soc. (3) **33** (1976), no. 2, 193–237, DOI 10.1112/plms/s3-33.2.193. MR0434947 ↑1480
- [27] The LMFDB Collaboration, *The L-functions and Modular Forms Database*, <http://www.lmfdb.org> 1461, 1465
- [28] Á. Lozano-Robledo, *On the field of definition of p -torsion points on elliptic curves over the rationals*, Math. Ann. **357** (2013), no. 1, 279–305, DOI 10.1007/s00208-013-0906-5. MR3084348 ↑1459, 1467, 1468, 1469, 1470, 1481
- [29] B. Mazur, *Modular curves and the Eisenstein ideal*, Inst. Hautes Études Sci. Publ. Math. **47** (1977), 33–186 (1978). With an appendix by Mazur and M. Rapoport. MR488287 ↑1458, 1460
- [30] B. Mazur, *Rational isogenies of prime degree (with an appendix by D. Goldfeld)*, Invent. Math. **44** (1978), no. 2, 129–162, DOI 10.1007/BF01390348. MR482230 ↑1459, 1461
- [31] L. Merel, *Bornes pour la torsion des courbes elliptiques sur les corps de nombres* (French), Invent. Math. **124** (1996), no. 1-3, 437–449, DOI 10.1007/s002220050059. MR1369424 ↑1458
- [32] F. Najman, *Torsion of rational elliptic curves over cubic fields and sporadic points on $X_1(n)$* , Math. Res. Lett. **23** (2016), no. 1, 245–272, DOI 10.4310/MRL.2016.v23.n1.a12. MR3512885 ↑1458, 1459, 1460, 1475
- [33] J. Rouse and D. Zureick-Brown, *Elliptic curves over \mathbb{Q} and 2-adic images of Galois*, Res. Number Theory **1** (2015), Art. 12, 34, DOI 10.1007/s40993-015-0013-7. MR3500996 ↑1478, 1480, 1481, 1482
- [34] J.-P. Serre, *Propriétés galoisiennes des points d'ordre fini des courbes elliptiques* (French), Invent. Math. **15** (1972), no. 4, 259–331, DOI 10.1007/BF01405086. MR0387283 ↑1461
- [35] J.-P. Serre, *Quelques applications du théorème de densité de Chebotarev* (French), Inst. Hautes Études Sci. Publ. Math. **54** (1981), 323–401. MR644559 ↑1462
- [36] A. V. Sutherland, *Computing images of Galois representations attached to elliptic curves*, Forum Math. Sigma **4** (2016), e4, 79, DOI 10.1017/fms.2015.33. MR3482279 ↑1461, 1462, 1482
- [37] D. Zywina, *On the possible images of the mod ℓ representations associated to elliptic curves over \mathbb{Q}* , arXiv:1508.07660. 1461, 1462, 1467, 1472, 1474, 1481

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, MADRID, SPAIN

Email address: enrique.gonzalez.jimenez@uam.es

URL: <http://matematicas.uam.es/~enrique.gonzalez.jimenez>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, 10000 ZAGREB, CROATIA

Email address: fnajman@math.hr

URL: <http://web.math.pmf.unizg.hr/~fnajman/>