

LEBESGUE-TYPE INEQUALITIES FOR QUASI-GREEDY BASES

EUGENIO HERNÁNDEZ

ABSTRACT. We show that for quasi-greedy bases in real Banach spaces the error of the thresholding greedy algorithm of order N is bounded by the best N -term error of approximation times a constant which depends on the democracy functions and the quasi-greedy constant of the basis.

1. INTRODUCTION

Let $(\mathbb{B}, \|\cdot\|_{\mathbb{B}})$ be a real Banach space with a countable seminormalized basis $\mathcal{B} = \{e_k : k \in \mathbb{N}\}$. Let $\Sigma_N, N = 1, 2, 3, \dots$ be the set of all $y \in \mathbb{B}$ with at most N non-null coefficients in the unique basis representation. For $x \in \mathbb{B}$, the N -term error of approximation with respect to \mathcal{B} is

$$\sigma_N(x) = \sigma_N(x; \mathcal{B}, \mathbb{B}) := \inf_{y \in \Sigma_N} \|x - y\|_{\mathbb{B}}, \quad N = 1, 2, 3, \dots$$

Given $x = \sum_{k \in \mathbb{N}} a_k(x) e_k \in \mathbb{B}$, let π denote any bijection of \mathbb{N} such that

$$|a_{\pi(k)}(x)| \geq |a_{\pi(k+1)}(x)| \quad \text{for all } k \in \mathbb{N}. \quad (1.1)$$

The **thresholding greedy algorithm of order N** (TGA) is defined by

$$G_N(x) = G_N^{\pi}(x; \mathcal{B}, \mathbb{B}) := \sum_{k=1}^N a_{\pi(k)}(x) e_{\pi(k)}.$$

It is not always true that $G_N(x) \rightarrow x$ (in \mathbb{B}) as $N \rightarrow \infty$. A basis \mathcal{B} is called **quasi-greedy** if $G_N(x) \rightarrow x$ (in \mathbb{B}) as $N \rightarrow \infty$ for all $x \in \mathbb{B}$. It turns out that this is equivalent (see Theorem 1 in [9]) to the existence of some constant C such that

$$\sup_N \|G_N(x)\|_{\mathbb{B}} \leq C \|x\|_{\mathbb{B}} \quad \text{for all } x \in \mathbb{B}. \quad (1.2)$$

It is convenient to define the **quasi-greedy constant K** to be the least constant such that

$$\|G_N(x)\|_{\mathbb{B}} \leq K \|x\|_{\mathbb{B}} \quad \text{and} \quad \|x - G_N(x)\|_{\mathbb{B}} \leq K \|x\|_{\mathbb{B}}, \quad x \in \mathbb{B}.$$

Given a basis \mathcal{B} in a Banach space \mathbb{B} a **Lebesgue-type inequality** is an inequality of the form

$$\|x - G_N(x)\|_{\mathbb{B}} \leq C v(N) \sigma_N(x), \quad x \in \mathbb{B},$$

where $v(N)$ is a nondecreasing function of N . For a survey of Lebesgue-type inequalities see [6] and the references given there.

Date: November 2, 2011.

2010 Mathematics Subject Classification. 41A65, 41A46, 41A17.

Key words and phrases. Lebesgue-type inequalities, thresholding greedy algorithm, quasi-greedy bases, democracy functions.

Research supported by Grant MTM2010-16518 (Spain).

The purpose of this note is to find Lebesgue-type inequalities for quasi-greedy basis in a Banach space.

For a seminormalized collection $\mathcal{B} = \{u_k\}_{k \in \mathbb{N}}$ in a Banach space \mathbb{B} the following quantities are defined:

$$h_r(N) = \sup_{|\Gamma|=N} \left\| \sum_{k \in \Gamma} u_k \right\|, \quad h_l(N) = \inf_{|\Gamma|=N} \left\| \sum_{k \in \Gamma} u_k \right\|,$$

and

$$\mu(N) = \sup_{1 \leq k \leq N} \frac{h_r(k)}{h_l(k)}, \quad N = 1, 2, 3, \dots \quad (1.3)$$

These functions are implicit in earlier works on N -term approximation and explicitly defined in [4]. The function $\mu(N)$ is defined in [9]. The functions h_r and h_l are called right and left democracy functions of \mathcal{B} (see [2] and [3]).

Our main result is the following:

Theorem 1.1. *Let $\mathcal{B} = \{e_k\}_{k=1}^\infty$ be a quasi-greedy basis in a real Banach space \mathbb{B} , and let K be the quasi-greedy constant of \mathcal{B} . Then for all $N = 1, 2, 3, \dots$ and all $x \in \mathbb{B}$,*

$$\|x - G_N(x)\|_{\mathbb{B}} \lesssim 8K^5 \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right) \sigma_N(x).$$

Before proving Theorem 1.1 we make some remarks about the function

$$v(N) := \sum_{k=1}^N \mu(k) \frac{1}{k}.$$

Obviously, μ is increasing so that $v(N) \lesssim \mu(N) \log N$. In some cases this inequality is an equivalence. For example if $\mu(N) \approx C$ (that is \mathcal{B} is democratic) then $v(N) \approx \log N$. In other cases the inequality can be improved. It can be proved that if \mathcal{B} is quasi-greedy, μ is doubling, that is there exists a constant $D \geq 1$ such that $\mu(2k) \leq D\mu(k)$, $k \in \mathbb{N}$ (see the Appendix). Under this condition it is not difficult to prove that

$$v(N) \approx \sum_{k=1}^{\log_2 N} \mu(2^k).$$

Moreover, if we assume that μ has a positive dilation index, that is $\mu \in \mathbb{W}_+$ in the terminology of [3], by Lemma 2.1 in [3] we have

$$v(N) \lesssim \mu(2^{\log_2 N}) = \mu(N),$$

so that in this situation we do not need the $\log N$ factor.

We prove Theorem 1.1 in Section 2. Section 3 contains some comments and open questions.

2. PROOF OF THEOREM 1.1

We need the following result from [1]: let $\mathcal{B} = \{e_k\}_{k=1}^\infty$ be a quasi-greedy bases with constant K in a real Banach space. For any finite set $\Gamma \subset \mathbb{N}$ and any real numbers $\{a_k\}_{k \in \Gamma}$ we have

$$\frac{1}{4K^2}(\min_{\Gamma} |a_k|) \left\| \sum_{k \in \Gamma} e_k \right\|_{\mathbb{B}} \leq \left\| \sum_{k \in \Gamma} a_k e_k \right\|_{\mathbb{B}} \leq (2K)(\max_{\Gamma} |a_k|) \left\| \sum_{k \in \Gamma} e_k \right\|_{\mathbb{B}} \quad (2.1)$$

(see Lemma 2.1 and Lemma 2.2 in [1]).

Lemma 2.1. *Let \mathcal{B} and \mathbb{B} as in Theorem 1.1. Suppose that there exists $C_1 > 0$ such that for all $\Gamma \subset \mathbb{N}$ finite*

$$\left\| \sum_{k \in \Gamma} e_k \right\|_{\mathbb{B}} \leq C_1 \eta(|\Gamma|) \quad (2.2)$$

for some η increasing and doubling (for example, $\eta(N) = h_r(N)$ and $C_1 = 1$). Then there exists $C = C_\eta$ such that for any $x = \sum_{k \in \mathbb{N}} a_k(x) e_k \in \mathbb{B}$

$$\|x\|_{\mathbb{B}} \leq 2K C_\eta \sum_{k=1}^{\infty} a_k^*(x) \eta(k) \frac{1}{k}, \quad (2.3)$$

where $\{a_k^*(x)\}$ is a decreasing rearrangement of $\{|a_k(x)|\}$ as in (1.1).

Proof. Let π be a bijection of \mathbb{N} that gives $\{a_k^*(x)\}$, that is $\{a_k^*(x)\} = \{a_{\pi_k}(x)\}$. Since \mathcal{B} is quasi-greedy

$$\lim_{N \rightarrow \infty} \sum_{k=1}^N a_{\pi_k}(x) e_{\pi_k} \rightarrow x \text{ (convergence in } \mathbb{B}\text{)}.$$

Thus

$$\begin{aligned} \|x\|_{\mathbb{B}} &= \left\| \sum_{k=1}^{\infty} a_{\pi(k)}(x) e_{\pi(k)} \right\|_{\mathbb{B}} = \left\| \sum_{j=0}^{\infty} \sum_{2^j \leq k < 2^{j+1}} a_{\pi(k)}(x) e_{\pi(k)} \right\|_{\mathbb{B}} \\ &\leq \sum_{j=0}^{\infty} \left\| \sum_{2^j \leq k < 2^{j+1}} a_{\pi(k)}(x) e_{\pi(k)} \right\|_{\mathbb{B}}. \end{aligned}$$

We now use first the right hand side inequality of (2.1) and then condition (2.2) to deduce

$$\begin{aligned} \|x\|_{\mathbb{B}} &= \sum_{j=0}^{\infty} (2K) |a_{\pi(2^j)}(x)| \left\| \sum_{2^j \leq k < 2^{j+1}} e_{\pi(k)} \right\|_{\mathbb{B}} \leq (2K) C_1 \sum_{j=0}^{\infty} |a_{\pi(2^j)}(x)| \eta(2^j) \\ &= (2K) C_1 \sum_{j=0}^{\infty} a_{2^j}^*(x) \eta(2^j). \end{aligned}$$

Inequality (2.3) follows since η is doubling and increasing. \square

Lemma 2.2. *Let \mathcal{B} and \mathbb{B} as in Theorem 1.1. Suppose that there exists $C_2 > 0$ such that for all $\Gamma \subset \mathbb{N}$ finite*

$$\frac{1}{C_2} \eta(|\Gamma|) \leq \left\| \sum_{k \in \Gamma} e_k \right\|_{\mathbb{B}} \quad (2.4)$$

for some function η (for example, $\eta(N) = h_l(N)$ and $C_2 = 1$). Then for any $x = \sum_{k \in \mathbb{N}} a_k(x) e_k \in \mathbb{B}$

$$\left[\sup_{\Gamma} a_k^*(x) \eta(k) \right] \leq C_2 (4K^3) \|x\|_{\mathbb{B}}, \quad (2.5)$$

where $\{a_k^*(x)\}$ is a decreasing rearrangement of $\{|a_k(x)|\}$ as in (1.1).

Proof. Let π be as in the proof of Lemma 2.1. For any $k \in \mathbb{N}$ we use condition (2.4) and then the left hand side inequality of (2.1) to obtain

$$|a_{\pi(k)}(x)|\eta(k) \leq C_2 |a_{\pi(k)}(x)| \left\| \sum_{j=1}^k e_{\pi(j)} \right\| \leq C_2 (4K^2) \left\| \sum_{j=1}^k a_{\pi(j)} e_{\pi(j)} \right\|.$$

We use (1.2) to deduce $|a_{\pi(k)}(x)|\eta(k) \leq C_2 (4K^3) \|x\|_{\mathbb{B}}$. The result follows by taking the supremum on $k \in \Gamma$. \square

For $\Gamma \subset \mathbb{N}$ and $x = \sum_{k \in \mathbb{N}} a_k(x) e_k \in \mathbb{B}$ define the **projection operator over Γ** as

$$S_{\Gamma}(x) := \sum_{k \in \Gamma} a_k(x) e_k.$$

Lemma 2.3. *Let \mathcal{B} and \mathbb{B} as in Theorem 1.1. For $\Gamma \subset \mathbb{N}$ finite*

$$\|S_{\Gamma}(x)\|_{\mathbb{B}} \lesssim (8K^4) \left(\sum_{k=1}^{|\Gamma|} \mu(k) \frac{1}{k} \right) \|x\|_{\mathbb{B}}.$$

Proof. Apply Lemma 2.1 with $\eta(N) = h_r(N)$ to obtain

$$\begin{aligned} \|S_{\Gamma}(x)\|_{\mathbb{B}} &= \left\| \sum_{k \in \Gamma} a_k(x) e_k \right\|_{\mathbb{B}} \lesssim (2K) \sum_{k=1}^{|\Gamma|} a_k^*(x) h_r(k) \frac{1}{k} \\ &\leq (2K) \sum_{k=1}^{|\Gamma|} a_k^*(x) \frac{h_r(k)}{h_l(k)} h_l(k) \frac{1}{k} \\ &\leq (2K) [\sup_k a_k^*(x) h_l(k)] \sum_{k=1}^{|\Gamma|} \mu(k) \frac{1}{k}. \end{aligned}$$

Use now Lemma (2.2) with $\eta(k) = h_l(k)$ to deduce the result. \square

We now prove Theorem 1.1. The proof follows arguments used in [4], [7] and [8] that were presented by V. N. Temlyakov at the *Concentration week on greedy algorithms in Banach spaces and compressed sensing* held on July 18-22 at Texas A&M University.

Take $\epsilon > 0$ and $N = 1, 2, 3, \dots$. Choose $p_N(x) = \sum_{k \in P} b_k e_k$ with $|P| = N$ such that

$$\|x - p_N(x)\|_{\mathbb{B}} \leq \sigma_N(x) + \epsilon. \quad (2.6)$$

Let Γ be the set of indices picked by the thresholding greedy algorithm after N iterations, that is

$$G_N(x) = \sum_{k \in \Gamma} a_k(x) e_k, \quad |\Gamma| = N.$$

We have, from Lemma 2.3 and (2.6)

$$\begin{aligned} \|x - G_N(x)\|_{\mathbb{B}} &\leq \|x - p_N(x)\|_{\mathbb{B}} + \|p_N(x) - S_P(x)\|_{\mathbb{B}} + \|S_P(x) - S_{\Gamma}(x)\|_{\mathbb{B}} \\ &= \|x - p_N(x)\|_{\mathbb{B}} + \|S_P(x - p_N(x))\|_{\mathbb{B}} + \|S_P(x) - S_{\Gamma}(x)\|_{\mathbb{B}} \end{aligned}$$

$$\begin{aligned}
&\lesssim \left[1 + 8K^4 \left(\sum_{k=1}^{|\Gamma|} \mu(k) \frac{1}{k} \right)\right] \|x - p_N(x)\|_{\mathbb{B}} + \|S_{P \setminus \Gamma}(x) - S_{\Gamma \setminus P}(x)\|_{\mathbb{B}} \\
&\leq \left[1 + 8K^4 \left(\sum_{k=1}^{|\Gamma|} \mu(k) \frac{1}{k} \right)\right] (\sigma_N(x) + \epsilon) + \|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}} + \|S_{\Gamma \setminus P}(x)\|_{\mathbb{B}}. \tag{2.7}
\end{aligned}$$

It is not difficult to find an upper bound for $\|S_{\Gamma \setminus P}(x)\|_{\mathbb{B}}$. Since $p_N(x)$ is supported in P we have $S_{\Gamma \setminus P}(p_N(x)) = 0$. By Lemma 2.3

$$\begin{aligned}
\|S_{\Gamma \setminus P}(x)\|_{\mathbb{B}} &= \|S_{\Gamma \setminus P}(x - p_N(x))\|_{\mathbb{B}} \lesssim (8K^4) \left(\sum_{k=1}^{|\Gamma \setminus P|} \mu(k) \frac{1}{k} \right) \|x - p_N(x)\|_{\mathbb{B}} \\
&\leq (8K^4) \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right) [\sigma_N(x) + \epsilon]. \tag{2.8}
\end{aligned}$$

The bound for $\|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}}$ is more delicate. Use Lemma 2.1 with $\eta(N) = h_r(N)$ to write

$$\|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}} = \left\| \sum_{k \in P \setminus \Gamma} a_k(x) e_k \right\|_{\mathbb{B}} \leq (2K) \sum_{k=1}^{|P \setminus \Gamma|} a_k^*(S_{P \setminus \Gamma}(x)) h_r(k) \frac{1}{k}.$$

If $k \in \Gamma \setminus P$ and $s \in P \setminus \Gamma$ we have $a_s^*(S_{P \setminus \Gamma}(x)) \leq a_k^*(S_{\Gamma \setminus P}(x))$ by construction of the thresholding greedy algorithm since $\min_{\Gamma} |a_k(x)| \geq \max_{N \setminus \Gamma} |a_k(x)|$. Also, since $|P| = N = |\Gamma|$ we have $|P \setminus \Gamma| = |\Gamma \setminus P|$. Thus

$$\|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}} \leq (2K) \sum_{k=1}^{|\Gamma \setminus P|} a_k^*(S_{\Gamma \setminus P}(x)) h_r(k) \frac{1}{k}.$$

We use that $S_{\Gamma \setminus P}(p_N(x)) = 0$ and Lemma 2.2 with $\eta = h_l$ to obtain

$$\begin{aligned}
\|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}} &\lesssim (2K) \sum_{k=1}^{|\Gamma \setminus P|} a_k^*(S_{\Gamma \setminus P}(x - p_N(x))) h_r(k) \frac{1}{k} \\
&= (2K) \sum_{k=1}^{|\Gamma \setminus P|} a_k^*(G_{|\Gamma \setminus P|}(x - p_N(x))) \frac{h_r(k)}{h_l(k)} h_l(k) \frac{1}{k} \\
&\leq (2K) [\sup_k a_k^*(G_{|\Gamma \setminus P|}(x - p_N(x))) h_l(k)] \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right) \\
&\leq (2K)(4K^3) \|G_{|\Gamma \setminus P|}(x - p_N(x))\|_{\mathbb{B}} \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right).
\end{aligned}$$

We use now that \mathcal{B} is a quasi-greedy basis to write

$$\begin{aligned}
\|S_{P \setminus \Gamma}(x)\|_{\mathbb{B}} &\lesssim (8K^5) \|x - p_N(x)\|_{\mathbb{B}} \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right) \\
&\leq (8K^5) (\sigma_N(x) + \epsilon) \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right). \tag{2.9}
\end{aligned}$$

Replacing (2.8) and (2.9) in (2.7), and letting $\epsilon \rightarrow 0$ we obtain the result stated in Theorem 1.1.

3. COMMENTS AND QUESTIONS

3.1. Let \mathcal{B} be a seminormalized quasi-greedy basis in a real Hilbert space \mathbb{H} . Since $\mathcal{B} = \{e_k\}_{k=1}^\infty$ is unconditional for constant coefficients (see Proposition 2 in [9]) it follows from Kintchine's inequality that

$$\left\| \sum_{k \in \Gamma} e_k \right\|_{\mathbb{H}} \approx \sqrt{|\Gamma|}.$$

Thus, in this case we can take $\eta(N) = N^{1/2}$ in Lemma 2.1 and Lemma 2.2, giving us Theorem 3 from [9].

3.2. Let $\mathcal{B} = \{e_k\}_{k=1}^\infty$ be a quasi-greedy basis in $L^p(\mathbb{T}^d)$. If $2 \leq p < \infty$ the space $L^p(\mathbb{T}^d)$ has type 2 and cotype p . Thus

$$C'_p |\Gamma|^{1/p} \leq \left\| \sum_{\Gamma} e_k \right\| \leq C_p |\Gamma|^{1/2}, \quad \Gamma \subset \mathbb{N}. \quad (3.1)$$

Taking $\eta(N) \approx N^{1/2}$ in Lemma 2.1 and $\eta(N) \approx N^{1/p}$ in Lemma 2.2 we obtain Theorem 11 from [7] for the case $2 \leq p < \infty$.

The case $1 < p \leq 2$ of Theorem 11 from [7] is obtained by observing that for this range of p 's the space $L^p(\mathbb{T}^d)$ has type p and cotype 2, so that

$$C'_p |\Gamma|^{1/2} \leq \left\| \sum_{\Gamma} e_k \right\| \leq C_p |\Gamma|^{1/p}, \quad \Gamma \subset \mathbb{N}. \quad (3.2)$$

3.3. The proofs of Lemmata 2.1 and 2.2 follow the pattern of the proofs of 1. \Rightarrow 2. in Theorem 3.1 and Theorem 4.2 from [3] for the limiting case " $\alpha = 0$ ".

3.4. As in [9] write, for $N = 1, 2, 3, \dots$

$$e_N(\mathbb{B}) = e_N := \sup_{x \in \mathbb{B}} \frac{\|x - G_n(x)\|_{\mathbb{B}}}{\sigma_N(x)}, \quad \left(\frac{0}{0} = 1\right).$$

Theorem 1.1 shows that for a quasi-greedy basis in a real Banach space

$$e_N \leq C \left(\sum_{k=1}^N \mu(k) \frac{1}{k} \right) \lesssim \mu(N) \log N, \quad N \in \mathbb{N}.$$

For greedy bases, Theorem 4 from [9] shows that

$$e_N \approx \mu(N), \quad N \in \mathbb{N}. \quad (3.3)$$

The same argument that proves (3.3) can be used to prove the following result: for a quasi-greedy basis \mathcal{B} in a real Banach space \mathbb{B}

$$\tilde{e}_N \approx \mu(N), \quad N \in \mathbb{N}. \quad (3.4)$$

were

$$\tilde{e}_N(\mathbb{B}) = \tilde{e}_N := \sup_{x \in \mathbb{B}} \frac{\|x - G_n(x)\|_{\mathbb{B}}}{\tilde{\sigma}_N(x)}, \quad \left(\frac{0}{0} = 1\right).$$

and

$$\tilde{\sigma}_N(x) = \tilde{\sigma}_N(x; \mathcal{B}, \mathbb{B}) := \inf \left\{ \left\| x - \sum_{k \in \Gamma} a_k(x) e_k \right\|_{\mathbb{B}} : |\Gamma| \leq N \right\}$$

is the *expansional* best approximation to $x = \sum_{k \in \mathbb{N}} a_k(x) e_k \in \mathbb{B}$.

Since $\sigma_N(x) \leq \tilde{\sigma}_N(x)$, for a quasi-greedy basis we have by (3.4) and Theorem 1.1

$$\mu(N) \lesssim \tilde{e}_N(\mathcal{B}) \leq e_N(\mathcal{B}) \lesssim \mu(N) \log N. \quad (3.5)$$

By the comments that follow the statement of Theorem 1.1 if μ has positive dilation index, $\mu(N) \lesssim \tilde{e}_N(\mathcal{B}) \leq e_N(\mathcal{B}) \lesssim \mu(N)$.

QUESTION 1. Is the inequality on the right hand side of (3.5) sharp? That is, is it possible to find a quasi-greedy basis \mathcal{B} such that $e_N(\mathcal{B}) \approx \mu(N) \log N$?

QUESTION 2. Is it true that for a quasi-greedy basis $\tilde{\sigma}_N(x) \lesssim \sigma_N(x) \log N$? If the answer is "yes" then by (3.4) we will have $e_N(\mathcal{B}) \lesssim \tilde{e}_N(\mathcal{B}) \log N \lesssim \mu(N) \log N$, given another proof of the right hand side of (3.5).

3.5. For a quasi-greedy basis $\mathcal{B} = \{e_k\}_{k=1}^{\infty}$ in $L^p(\mathbb{T}^d)$, inequalities 3.1 and 3.2 (or type and cotype properties of $L^p(\mathbb{T}^d)$) show that $\mu(N) \lesssim N^{|\frac{1}{p}-\frac{1}{2}|}$. By the comments that follow the statement of Theorem 1.1, if $p \neq 2$ and $1 < p < \infty$, $e_N(\mathbb{B}) \lesssim N^{|\frac{1}{p}-\frac{1}{2}|}$, proving Theorem 1.1 from [8]. (Notice that $w(N) := N^{|\frac{1}{p}-\frac{1}{2}|}$ has positive dilation index if $p \neq 2$.) For $p = 2$ we have $e_N(\mathcal{B}) \lesssim \log N$ by Theorem 1.1.

Consider now the trigonometric system $\mathcal{T}^d = \{e^{ikx} : k \in \mathbb{Z}^d\}$ in $L^p(\mathbb{T}^d)$, $1 \leq p \leq \infty$ (here $L^\infty(\mathbb{T}^d)$ is $C(\mathbb{T}^d)$, the set of continuous functions in \mathbb{T}^d). It is proved in [5] (Theorem 2.1) that

$$e_N(\mathcal{T}^d, L^p(\mathbb{T}^d)) \lesssim N^{|\frac{1}{p}-\frac{1}{2}|}, \quad 1 \leq p \leq \infty.$$

QUESTION 3. (Asked by V. N. Temlyakov at the *Concentration week on greedy algorithms in Banach spaces and compressed sensing* held on July 18-22 at Texas A&M University.)

a) Characterize those systems \mathcal{B} in $L^p(\mathbb{T}^d)$, $1 \leq p \leq \infty$, such that $e_N(\mathcal{T}^d, L^p(\mathbb{T}^d)) \lesssim N^{|\frac{1}{p}-\frac{1}{2}|}$, $N \in \mathbb{N}$. Notice that if $1 < p \neq 2 < \infty$, the characterization must be satisfy by \mathcal{T}^d as well as any quasi-greedy basis.

More generally,

b) Let $v(N)$ be an increasing function of N . Characterize those systems \mathcal{B} in a Banach space \mathbb{B} for which $e_N(\mathcal{B}, \mathbb{B}) \lesssim v(N)$.

4. APPENDIX

Lemma 4.1. *If \mathcal{B} is a quasi-greedy basis in a Banach space \mathbb{B} , the function μ defined in (1.3) is doubling.*

Proof. It is proved in [9] and [1] that for a quasi-greedy basis $\mathbb{B} = \{e_k\}_{k=1}^{\infty}$ with quasi-greedy constant K , if $B \subset A \subset \mathbb{N}$ (finite sets) then

$$\left\| \sum_{k \in B} e_k \right\|_{\mathbb{B}} \leq K \left\| \sum_{k \in A} e_k \right\|_{\mathbb{B}}. \quad (4.1)$$

We have to prove that $\mu(2N) \leq D\mu(N)$ for some D independent of N . Since $\mu(2N)$ is defined as a supremum over the finite set $1 \leq k \leq 2N$, there exists $k_0 \leq 2N$ such

that $\mu(2N) = h_r(k_0)/h_l(k_0)$. Notice that h_r is doubling with doubling constant 2 by the triangle inequality.

Suppose first that $k_0 = 2s \leq 2N$ is even. From (4.1) we deduce $h_l(s) \leq Kh_l(2s)$. Hence

$$\mu(2N) = \frac{h_r(2s)}{h_l(2s)} \leq (2K) \frac{h_r(s)}{h_l(s)} \leq (2K)\mu(N)$$

since $s \leq N$.

Assume now that $k_0 = 2s + 1$ is odd. Since $2s + 1 = k_0 \leq 2N$ we deduce $s \leq N - \frac{1}{2}$, and since s is an integer $s \leq N - 1$. From (4.1) we deduce $h_r(2s + 1) \leq Kh_r(2s + 2)$ and $h_l(s + 1) \leq Kh_l(2s + 1)$. Hence

$$\mu(2N) = \frac{h_r(2s + 1)}{h_l(2s + 1)} \leq K^2 \frac{h_r(2s + 2)}{h_l(s + 1)} \leq 2K^2 \frac{h_r(s + 1)}{h_l(s + 1)} \leq (2K^2)\mu(N)$$

since $s + 1 \leq N$. □

Acknowledgements. This work started when the author participated in the *Concentration week on greedy algorithms in Banach spaces and compressed sensing* held on July 18-22 at Texas A&M University. I would like to express my gratitude to the Organizing Committee for the invitation to participate in this meeting.

REFERENCES

- [1] S.J. DILWORTH, N.J. KALTON, D. KUTZAROVA, V.N. TEMLYAKOV, *The Thresholding Greedy Algorithm, Greedy Bases, and Duality*, Constr. Approx., 19, (2003), 575–597.
- [2] G. GARRIGÓS, E. HERNÁNDEZ, J.M. MARTELL, *Wavelets, Orlicz spaces and greedy bases*, Appl. Compt. Harmon. Anal., (24): (2008), 70–93.
- [3] G. GARRIGÓS, E. HERNÁNDEZ, M. DE NATIVIDADE, *Democracy functions and optimal embeddings for approximation spaces*, Adv. in Comp. Math. 2011 (Accepted)
- [4] A. KAMONT, V.N. TEMLYAKOV, *Greedy approximation and the multivariate Haar system*, Studia Math, 161 (3), (2004), 199–223.
- [5] V. N. TEMLYAKOV, *Greedy algorithm and n-term trigonometric approximation*, Const.Approx., 14, (1998), 569–587.
- [6] V. N. TEMLYAKOV, *Greedy approximation*, Acta Numerica (2008), 235–409.
- [7] V. N. TEMLYAKOV, M. YANG, P. YE, *Greedy approximation with regard to non-greedy bases*, Adv. in Comp. Math., 34, (2011), 219–337.
- [8] V. N. TEMLYAKOV, M. YANG, P. YE, *Lebesgue-type inequalities for greedy approximation with respect to quasi-greedy bases*, Preprint.
- [9] P. WOJSTASZCZYK, *Greedy Algorithm for General Biorthogonal Systems*, Journal of Approximation Theory, 107, (2000), 293–314.

EUGENIO, HERNÁNDEZ, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049, MADRID, SPAIN

E-mail address: eugenio.hernandez@uam.es