

Total Variation Flow and Sign Fast Diffusion in one dimension

Matteo Bonforte^a and Alessio Figalli^b

August 18, 2011

Abstract

We consider the dynamics of the Total Variation Flow (TVF) $u_t = \operatorname{div}(Du/|Du|)$ and of the Sign Fast Diffusion Equation (SFDE) $u_t = \Delta \operatorname{sign}(u)$ in one spatial dimension. We find the explicit dynamic and sharp asymptotic behaviour for the TVF, and we deduce the one for the SFDE by an explicit correspondence between the two equations.

Keywords. Fast Diffusion, Total Variation Flow, asymptotic, extinction profile, convergence rates, extinction time.

Mathematics Subject Classification. 35K55, 35K59, 35K67, 35K92, 35B40.

(a) Departamento de Matemáticas, Universidad Autónoma de Madrid, Campus de Cantoblanco, 28049 Madrid, Spain.

E-mail: matteo.bonforte@uam.es. Web-page: <http://www.uam.es/matteo.bonforte>

(b) Department of Mathematics, The University of Texas at Austin, 1 University Station C1200, Austin, TX 78712-1082, USA.

E-mail: figalli@math.utexas.edu. Web-page: <http://www.ma.utexas.edu/users/figalli/>

1 Introduction

The Total Variation Flow (TVF) and the Sign-Fast Diffusion Equation (SFDE) are two degenerate parabolic equations arising respectively as the limit of the (parabolic) p -Laplacian as $p \rightarrow 1^+$ and of the fast diffusion equations as $m \rightarrow 0^+$. In one spatial dimension these two equations are strictly related, and our goal here is to describe their dynamics.

Let us first introduce the SFDE: consider the Fast Diffusion Equation (FDE)

$$\partial_t v = \Delta(v^m), \quad 0 < m < 1 \quad (1.1)$$

(by definition, $v^m = |v|^{m-1}v$). By letting $m \rightarrow 0^+$ one gets the SFDE

$$\partial_t v = \Delta(\text{sign}(v)).$$

To study the evolution of this equation in one dimension, we will exploit its relation with the TVF (also called 1-Laplacian, as it corresponds to the limit of the p -Laplacian when $p \rightarrow 1^+$): at least formally, if v solves the SFDE, then $u(x) := \int_0^x v$ solves the TVF

$$\partial_t u = \text{div} \left(\frac{Du}{|Du|} \right). \quad (1.2)$$

Of course this is purely formal, and we will need to justify it, see Section 3. The correspondence between solutions of the p -Laplacian type equations and solutions to fast diffusion type equations have been used since a long time, cf. [24] and more recently in [6], in the study of equations related to the fronts represented in image contour enhancement. In several dimension the correspondence between solutions of the p -Laplacian and of the FDE is less explicit and holds only for radial solutions, cf. [20]. Here our strategy is first to analyze the dynamic of the TVF in one dimension, and then use this to recover the behaviour of solutions of the SFDE.

The literature on the TVF is quite rich, and we suggest to the interested reader the monograph [5] as source of references for the existence, uniqueness, basic regularity and different concepts of solutions and their relations, together with estimates on the extinction time (see also the review paper [16] and references therein). The TVF has some interest in applications to noise reduction, cf. [23, 1, 5]. The asymptotic behaviour of the TVF is still an open problem in many aspects, even if partial results have appeared in [4, 3, 5, 7, 8, 9, 17]. However, at least in the simpler case of one spatial dimension, the results contained in the present paper exhibit an almost explicit dynamic and a sharp asymptotic behaviour. After the writing of this paper was essentially completed, we learned of a related work [19].

Plan of the paper.

- In section 2 we analyze the dynamic of the one-dimensional TVF. As a first step in this direction, we study the time discretized case, for which we find the explicit dynamics for “local step functions”, see Subsection 2.3 and 2.4. In Subsection 2.5 we pass to the continuous time case, and we find the explicit evolution under the TVF for a generic “local step functions”. Then, in Subsection 2.6, exploiting the stability of the TVF in L^p spaces and arguing by approximation we prove some basic but important properties of solutions to the TVF, such as the conservation and contractivity property of the local modulus of continuity (Theorem 2.8), and the explicit behaviour around maxima and minima.

In the case of nonnegative compactly supported initial data, we first prove an explicit formula for the loss of mass and extinction time for the associated solution to the TVF (Proposition 2.10). Next we study the asymptotic behaviour of such solutions: we analyze and classify the possible asymptotic profiles (that we shall call more properly “extinction profiles”, see Theorem 2.11), and we characterize the asymptotic behaviour of solution to the TVF near the extinction time as a function of the initial datum (see Theorem 2.15). To our knowledge this is the first completely explicit asymptotic result for very singular parabolic equations, even if it holds in only one spatial dimension. Finally, in Subsection 2.10 we prove the sharpness of the rate of convergence provided by Theorem 2.15.

- In Section 3 we dedicate our attention to the SFDE. First we rigorously show that a BV function solves the TVF if and only if its distributional derivative solves the SFDE, and then we exploit this to describe the dynamic of the SFDE. We conclude the paper with a discussion on the relation between the SFDE and the Logarithmic Fast Diffusion Equation (LFDE), which is another possible limiting equation of the Fast Diffusion Equation as $m \rightarrow 0^+$, see Subsection 3.2.

ACKNOWLEDGEMENTS: We warmly thank J. L. Vázquez for useful comments and discussions. A.F. was partially supported by the NSF grant DMS-0969962. M.B. has been partially funded by Project MTM2008-06326-C02-01 and Ramon y Cajal grant RYC-2008-03521 (Spain).

2 The 1-dimensional Total Variation Flow

In this part we deal with the one dimensional TVF. Before introducing the problem, we first recall briefly some notation and basic facts about BV functions for convenience of the reader.

2.1 Notations and basic facts about BV functions in one dimension

Here we recall some basic facts about one-dimensional BV functions, referring to [2, Section 3.2] for more details.

Consider an open connected interval $I \subseteq \mathbb{R}$. A function $u \in BV(I)$ if $u \in L^1_{loc}(I)$, its distributional derivative Du is a (signed) measure, and its total variation $|Du|$ has finite mass.

The distributional derivative Du can be decomposed as $Du = \partial_x u \, dx + D^s u$, where $\partial_x u = \nabla u$ is the absolutely continuous part of Du (with respect to the Lebesgue measure), and $D^s u$ is the singular part.

By Sobolev inequalities we have the inclusion $BV(\mathbb{R}) \subset L^\infty(\mathbb{R})$, and $BV(I) \cap L^1(I) \subset L^p(I)$ for any $1 \leq p \leq \infty$.

If $u \in BV(I)$, up to redefining the function in a set of measure zero, for every point $x \in I$ it always exists the left or right limit of u at a point x , which we denote by

$$u(x^\pm) := \lim_{y \rightarrow x^\pm} u(y).$$

Moreover, the limits above are equal up to a countable number of points. We will always assume to work with a “good representative”, so that the above property always holds (see [2, Theorem 3.28]).

2.2 The setting

Let us briefly recall the definition of strong solution to the TVF, in the form we will use it throughout this paper. For the moment, we do not specify any boundary condition (so the following discussion could be applied to the Cauchy problem in \mathbb{R} , as well as the Dirichlet or the Neumann problem on an interval).

A function $u \in L^\infty([0, \infty), BV(I)) \cap W_{loc}^{1,2}([0, \infty), L^2(I))$ is a strong solution of the TVF if there exists $z \in L_{loc}^2([0, \infty), W^{1,2}(I))$, with $\|z\|_\infty \leq 1$, such that

$$\partial_t u = \partial_x z \quad \text{on} \quad (0, \infty) \times I, \quad (2.1)$$

and

$$\int_0^T \int_I z(t, x) Du(t, x) dt dx = \int_0^T \int_I |Du(t, x)| dx dt \quad \forall T > 0.$$

Roughly speaking, the above condition says that $z = Du/|Du|$. We refer to the book [5] for a more detailed discussion on the different concepts of solution to the TVF depending on the classes of initial data (entropy solutions, mild solutions, semigroup solution), and equivalence among them.

Throughout the paper we will deal with non-negative initial data for the TVF, although many properties maybe extended to signed initial data.

2.3 The analysis of the time-discretized problem.

It is well known that the strong solution u of the TVF defined above is generated via Crandall-Liggett's Theorem, namely it is obtained as the limit of solutions of a time-discretized problem, formally given by the implicit Euler scheme

$$\frac{u(t_{i+1}) - u(t_i)}{t_{i+1} - t_i} = \partial_x \left(\frac{Du(t_{i+1})}{|Du(t_{i+1})|} \right).$$

We refer to the book [5] for a more complete and detailed discussion of these facts. The goal of this section is to understand the behaviour of the time-discretized solution both at continuity and at discontinuity points (see Propositions 2.2 and 2.3).

Let us fix a time step $h > 0$, set $t_0 = 0$, $t_{i+1} = t_i + h = (i+1)h$, and define $u_{ih}(x) := u(ih, x)$ so that $u_0(x) = u(0, x)$. The first step reads:

$$\frac{u_h(x) - u_0(x)}{h} = \partial_x z_h, \quad (2.2)$$

where $z_h \in L^\infty(I)$ satisfies $\|z_h\|_\infty \leq 1$, $z_h Du_h = |Du_h|$, and $\partial_x z \in L^2(I)$. Of course it suffices to understand the behavior of u_h starting from u_0 , as all the other steps will follow then by iteration.

We are going to prove the main properties of the time discretized solution, and to this end is useful to recall an equivalent definition for u_h :

$$u_h = \operatorname{argmin}[\Phi_h(u)], \quad \text{where} \quad \Phi_h(u) = \int_I |Du| + \frac{1}{2h} \int_I |u - u_0|^2 dx. \quad (2.3)$$

Indeed by strict convexity of the functional Φ_h , the minimizer is unique and is uniquely characterized by the Euler-Lagrange equation associated to Φ_h , which is exactly (2.2), that we shall rewrite in the form

$$u_h = u_0 + h \partial_x z_h. \quad (2.4)$$

Let us observe that the above construction does not need u_0 to be $L^2(I)$: if $u_0 \in BV(I)$ the above scheme still makes sense and provides a function u_h such that $u_h - u_0 \in L^2(I)$.

Next, we remark that since $u_0, u_h \in BV(I)$ also $\partial_x z_h \in BV(I) \subset L^\infty(I)$, which implies that z_h is Lipschitz and is differentiable outside a countable set of points. Define the (at most countable) set

$$N(z_h) := \left\{ x \in \mathbb{R} \mid \lim_{\varepsilon \rightarrow 0} \frac{z_h(x + \varepsilon) - z_h(x)}{\varepsilon} \text{ does not exist} \right\}. \quad (2.5)$$

Since $\partial_x z_h \in BV(I)$, it is continuous outside $N(z_h)$ (i.e. $\partial_x z_h \in C^0(\mathbb{R} \setminus N(z_h))$), and we have that $N(z_h)$ coincides with the set of discontinuity point of $u_h - u_0$.

Collecting all the information obtained so far, we can say that equation (2.4) is equivalent to

$$\begin{cases} h \partial_x z_h(x) = u_h(x) - u_0(x) & \text{for all } x \in \mathbb{R} \setminus N(z_h) \\ |z_h(x)| \leq 1, & \text{for all } x \in \mathbb{R} \\ z_h(x) = \pm 1, & \text{for } |Du_h| \text{ - a.e.} \end{cases} \quad (2.6)$$

The next lemmata will allow us to show the important fact that, on $\mathbb{R} \setminus N(z_h)$, u_h is locally constant whenever different from u_0 (see Proposition 2.2).

Lemma 2.1 *The following holds:*

$$|Du_h|(\{\partial_x z_h \neq 0\} \setminus N(z_h)) = |Du_h|(\{u_h \neq u_0\} \setminus N(z_h)) = 0. \quad (2.7)$$

Proof. The equality

$$\{\partial_x z_h \neq 0\} \setminus N(z_h) = \{u_h \neq u_0\} \setminus N(z_h) \quad (2.8)$$

easily follows by observing that $u_h - u_0 = h \partial_x z_h \in C^0(\mathbb{R} \setminus N(z_h))$. Moreover, since $z_h \cdot Du_h = |Du_h|$, we have

$$\int_{\mathbb{R}} (1 - |z_h(x)|) d|Du_h| = 0$$

which in particular implies

$$\int_{\{u_h \neq u_0\} \setminus N(z_h)} (1 - |z_h(x)|) d|Du_h| = 0. \quad (2.9)$$

Next we notice that since z_h is differentiable on $\mathbb{R} \setminus N(z_h)$ and $|z_h| \leq 1$, we have $\{z_h = \pm 1\} \cap \{\partial_x z_h \neq 0\} \cap (\mathbb{R} \setminus N(z_h)) = \emptyset$. Hence, thanks to (2.8) we deduce that $(1 - |z_h(x)|) > 0$ on the set $\{u_h \neq u_0\} \setminus N(z_h)$. Combining this information with (2.9) we obtain that $|Du_h|(\{u_h \neq u_0\} \setminus N(z_h)) = 0$, as desired. \square

We now use the above lemma to analyze the behavior of u_h near continuity points.

Proposition 2.2 (Behaviour near continuity points) *If u_h is different from u_0 at some common continuity point x , then it is constant in an open neighborhood of x .*

Proof. Let $x \in I$ be a continuity point both for u_0 and u_h , and assume that $u_h(x) > u_0(x)$ (the case $u_h(x) < u_0(x)$ being analogous). Then by (2.6) we deduce that $\partial_x z_h$ is continuous and strictly positive in an open neighborhood $I(x)$ of x , which together with (2.7) implies

$$|Du_h|(I(x)) = 0.$$

Hence u_h is constant on $I(x)$. \square

We now show that if $u_0 \in BV(I)$ has some discontinuity jump, then u_h can only have jumps at such points, and moreover the size of such jumps cannot increase.

Lemma 2.3 (Behaviour at discontinuity points) *Let $u_0 \in BV(I)$. Then, the following inequalities hold for any $x \in I$:*

$$\begin{aligned} \text{if } u_h(x^-) \leq u_h(x^+) \quad \text{then } u_0(x^-) \leq u_h(x^-) < u_h(x^+) \leq u_0(x^+) \\ \text{if } u_h(x^+) \leq u_h(x^-) \quad \text{then } u_0(x^+) \leq u_h(x^+) < u_h(x^-) \leq u_0(x^-). \end{aligned} \quad (2.10)$$

Moreover,

$$\begin{aligned} u_h(x^-) < u_h(x^+) \quad \text{implies } z_h(x) = 1 \\ u_h(x^-) > u_h(x^+) \quad \text{implies } z_h(x) = -1. \end{aligned} \quad (2.11)$$

Proof. Let $x \in I$ be a discontinuity point for u_h . Then

$$Du_h(x) = (u_h(x^+) - u_h(x^-))\delta_x, \quad (2.12)$$

where δ_x is the Dirac delta at $x \in I$. We first prove (2.11): recalling that $z_h \cdot Du_h = |Du_h|$, if $u_h(x^-) < u_h(x^+)$ then by (2.12) we get $z_h(x) = 1$. Analogously $u_h(x^-) > u_h(x^+)$ implies $z_h(x) = -1$.

Let us now show (2.10): assume first that $u_h(x^-) < u_h(x^+)$. Since $z_h(x) = 1$, x is a maximum point for z_h , thus $\partial_x z_h(x^-) \geq 0$ and $\partial_x z_h(x^+) \leq 0$. Using (2.6), this implies

$$\frac{u_h(x^+) - u_0(x^+)}{h} = \partial_x z_h(x^+) \leq 0 \leq \partial_x z_h(x^-) = \frac{u_h(x^-) - u_0(x^-)}{h}$$

which combined with our assumption $u_h(x^-) < u_h(x^+)$ gives

$$u_0(x^-) \leq u_h(x^-) < u_h(x^+) \leq u_0(x^+).$$

The case $u_h(x^-) > u_h(x^+)$ is analogous. \square

As an immediate corollary we get:

Corollary 2.4 (Local continuity) *Let $x \in I$. If u_0 is continuous at x , then u_h is continuous at x .*

This result shows that (at least at the discrete level) the TVF cannot create new discontinuities, and that continuous initial data produce continuous solutions (this is actually what we will prove in Theorem 2.8). Observe that this is a local property which does not depend on the boundary conditions (that we have not specified yet).

We conclude this section with the following estimates on the local loss of mass.

Lemma 2.5 (Local L^1 -estimates) *The following estimates hold for any interval $(a, b) \subseteq I$:*

$$\left| \int_a^b u_h(x) \, dx - \int_a^b u_0(x) \, dx \right| \leq 2h. \quad (2.13)$$

Proof. By equation (2.6) we have

$$z_h(b) - z_h(a) = \int_a^b \partial_x z_h(x) \, dx = \frac{1}{h} \left[\int_a^b u_h(x) \, dx - \int_a^b u_0(x) \, dx \right]. \quad (2.14)$$

Hence (2.13) follows from the bound $\|z_h\|_\infty \leq 1$. \square

2.4 The dynamics of local step functions I. The time discretized case

In this section we use the time discretization scheme to study the dynamics for initial data u_0 which coincide with a step function on some open interval I .

Let us point out that, if u_0 is exactly a step function, then one can give an explicit formula for its evolution (see Section 2.4.2 and [17]) by simply checking that it satisfies the equation. However, by studying the “time-discretized” evolution (and then letting $h \rightarrow 0$), one can see in a much more natural way the “locality” in the dynamic of the TVF. Moreover, our method shows how to deal with functions which do not belong to $L^2(I)$. Finally, our description give a good insight of the analysis of the discretized PDE, which may be useful for numerical purposes.

We would like to notice that it is important for the sequel that the time step $h > 0$ is sufficiently small with respect to the size of the jumps of the step function we are considering, as otherwise the discretized dynamics becomes more involved, as we shall show with an example at the end of this section.

2.4.1 Local evolution of a single step.

To give an insight on the way the discretized evolution behaves, we consider the case of *maximum steps*, whose behavior is made clear in Figure 1. Let us fix an interval $I = I_1 \cup I_2 \cup I_3$, and assume that $u_0 = \alpha_1 \chi_1 + \alpha_2 \chi_2 + \alpha_3 \chi_3$ on I , with $\alpha_2 > \max\{\alpha_1, \alpha_3\}$ and $\chi_k = \chi_{I_k}$ is the characteristic function of the open interval $I_k = (x_{k-1}, x_k)$. (Observe that we make no assumptions on u_0 outside I .) Fix $h > 0$ small (the smallness to be fixed), and consider the function u_h . We have:

- (a) u_h is constant on any interval I_k . This follows easily from by Proposition 2.2 and Corollary 2.4.
- (b) If h is small enough, then u_h jumps at the points x_1 and x_2 . Indeed, if by contradiction x_1 is a continuity point for u_h , u_h would equal to a constant $\bar{\alpha}$ on $I_1 \cup I_2$, with $\alpha_1 \leq \bar{\alpha} \leq \alpha_2$




Figure 1: *Dynamics of a maximum step.* This figure shows the dynamic only inside the interval $[x_0, x_3]$.

(see Lemma 2.3). However, by Lemma 2.5 we know that

$$|\alpha_1 - \bar{\alpha}| |I_1| = \left| \int_{I_1} [u_h(x) - u_0(x)] dx \right| \leq 2h$$

and

$$|\alpha_2 - \bar{\alpha}| |I_2| = \left| \int_{I_2} [u_h(x) - u_0(x)] dx \right| \leq 2h$$

which is impossible if

$$0 < h < \frac{|\alpha_1 - \alpha_2|}{2} \left(\frac{1}{|I_1|} + \frac{1}{|I_2|} \right)^{-1}. \quad (2.15)$$

In particular, u_h jumps both at x_1 and x_2 if the “simpler” condition

$$0 < h < |\alpha_1 - \alpha_2| \min \{|I_1|, |I_2|\} \quad (2.16)$$

holds.

(c) Applying Lemma 2.3 again (see equation (2.11)) we obtain

$$z_h(x_1) = 1 \quad \text{and} \quad z_h(x_2) = -1,$$

which combined with (2.14) gives

$$\frac{1}{h} \left[\int_{x_1}^{x_2} u_h(x) dx - \int_{x_1}^{x_2} u_0(x) dx \right] = -2.$$

Hence

$$u_h = \alpha_{2,h} := \alpha_2 - \frac{2h}{|I_2|} \quad \text{on } I_2.$$

(d) Combining all together we obtain that

$$u_0 = \alpha_1 \chi_1 + \alpha_2 \chi_2 + \alpha_3 \chi_3 \quad \text{on } I \quad \text{implies} \quad u_h = \alpha_{1,h} \chi_1 + \underbrace{\left(\alpha_2 - \frac{2h}{|I_2|} \right)}_{\alpha_{2,h}} \chi_2 + \alpha_{3,h} \chi_3 \quad \text{on } I, \quad (2.17)$$

where $\alpha_1 \leq \alpha_{1,h} \leq \alpha_{2,h}$, $\alpha_3 \leq \alpha_{3,h} \leq \alpha_{2,h}$ (see Lemma 2.3). The exact value of $\alpha_{1,h}$ and $\alpha_{3,h}$ depend on the behavior of u_0 outside I , but using (2.14) we can always estimate them:

$$|\alpha_{1,h} - \alpha_1| \leq \frac{2h}{|I_1|}, \quad |\alpha_{3,h} - \alpha_3| \leq \frac{2h}{|I_3|}. \quad (2.18)$$

(In some explicit cases where one knows that value of z at x_1 and x_3 , $\alpha_{1,h}$ and $\alpha_{3,h}$ can be explicitly computed using (2.14).)

Remark 2.6 It is important to observe that the value of $\alpha_{2,h}$ is *independent* of the value of $\alpha_{1,h}$ and $\alpha_{3,h}$, but only depends on the fact that $\alpha_1, \alpha_3 < \alpha_{2,h}$. In particular, thanks to (2.18), one can iterate the above construction: after ℓ steps we get

$$u_{\ell h} = \alpha_{1,\ell h} \chi_1 + \alpha_{2,\ell h} \chi_2 + \alpha_{3,\ell h} \chi_3 \quad \text{on } I, \quad \alpha_{2,\ell h} = \alpha_2 - \frac{2\ell h}{|I_2|}$$

holds as long as $\alpha_{1,(\ell-1)h}, \alpha_{3,(\ell-1)h} < \alpha_{2,\ell h}$, which for instance is the case (by iterating the estimate (2.18)) if

$$\ell h < |\alpha_1 - \alpha_2| \min\{|I_1|, |I_2|\} \quad \text{and} \quad \ell h < |\alpha_2 - \alpha_3| \min\{|I_2|, |I_3|\}.$$

In particular observe that in this analysis we never used that I_1 and I_3 are bounded intervals, so the above formulas also holds when $x_0 = -\infty$ and $x_3 = +\infty$.

2.4.2 Evolution of a general step function

The above analysis can be easily extended to the general N -step function: assume that

$$u_0 = \sum_{k=0}^{N+1} \alpha_k \chi_k \quad \text{on } I$$

where $\alpha_k \in \mathbb{R}$ for $k = 0, \dots, N+1$, and $\chi_k = \chi_{I_k}$ is the characteristic function of the open interval $I_k = (x_{k-1}, x_k)$ (also the values $x_0 = -\infty$ and $x_{N+1} = +\infty$ are allowed). Then, if

$$0 < \ell h < \min_{j=0, \dots, N} \left\{ |\alpha_j - \alpha_{j+1}| \min\{|I_j|, |I_{j+1}|\} \right\}, \quad (2.19)$$

the discrete solution after ℓ steps is given by

$$u_{\ell h} = \sum_{k=0}^{N+1} \alpha_{k,\ell h} \chi_k \quad \text{on } I,$$

where we are able to explicitly get the values of $\alpha_{k,\ell h}$ for $k = 1, \dots, N$, see Remark 2.6, and some information on $\alpha_{0,\ell h}$ and $\alpha_{N+1,\ell h}$: for $k = 1, \dots, N$

$$\alpha_{k,\ell h} = \begin{cases} \alpha_k, & \text{if } \alpha_{k-1} < \alpha_k < \alpha_{k+1} \text{ or if } \alpha_{k+1} < \alpha_k < \alpha_{k-1} \\ \alpha_k - \frac{2\ell h}{|I_k|}, & \text{if } \alpha_k > \max\{\alpha_{k-1}, \alpha_{k+1}\} \\ \alpha_k + \frac{2\ell h}{|I_k|}, & \text{if } \alpha_k < \min\{\alpha_{k-1}, \alpha_{k+1}\} \end{cases} \quad (2.20)$$

$$\alpha_{0,\ell h} \begin{cases} \geq \alpha_{0,(\ell-1)h}, & \text{if } \alpha_0 < \alpha_1 \\ \leq \alpha_{0,(\ell-1)h}, & \text{if } \alpha_0 > \alpha_1 \end{cases}$$

$$\alpha_{N+1,\ell h} \begin{cases} \geq \alpha_{N+1,(\ell-1)h}, & \text{if } \alpha_N > \alpha_{N+1} \\ \leq \alpha_{N+1,(\ell-1)h}, & \text{if } \alpha_N < \alpha_{N+1}. \end{cases}$$

A concluding remark on the smallness of the time step h . Since we want to describe the behaviour of the TVF, we are mainly interested in the limit $h \rightarrow 0$, which means that condition (2.19) is always fulfilled. Anyway it is interesting to observe that the dynamic becomes more complicated to understand for general values of h , since the “locality” property is lost. Figure 2 shows a situation when a maximum and a minimum disappear in one step (for this to happen, the area A has to be less than $2h$). Of course one can construct much more complicated examples. With this one, we can observe that the value of u_h inside $[x_1, x_2]$ depends on the values of u_0 on both $[x_1, x_2]$ and $[x_2, x_3]$.




Figure 2: Dynamics of a maximum step when h is not necessarily small. This figure shows the dynamic only inside the interval $[x_0, x_4]$.

2.5 The dynamics of local step functions II. The continuous time case

We now deduce the explicit evolution of strong solutions to the TVF introduced in Section 2.3 for the Cauchy problem on \mathbb{R} where the initial data coincide with a step function (see Remark 2.7 below for the analysis of initial value problems with boundary conditions on intervals). The dynamic of step functions will then be used in the next section to deduce, by approximation, qualitative properties for general solutions.

With the same notation as in Section 2.4.2, we consider the initial data given by

$$u_0(x) = \sum_{k=0}^{N+1} \alpha_k \chi_{I_k}(x) \quad \text{inside } I. \quad (2.21)$$

Then, for h small and $x \in I$, we define

$$u^h(t, x) := \left(\ell + 1 - \frac{t}{h} \right) u_{\ell h}(x) + \left(\frac{t}{h} - \ell \right) u_{(\ell+1)h}(x) \quad \text{for any } t \in [\ell h, (\ell+1)h].$$

Then it is immediate to check that, by (2.19), on the time interval $[0, t_1]$ with

$$t_1 < \min_{j=0, \dots, N} \left\{ |\alpha_j - \alpha_{j+1}| \min \{|I_j|, |I_{j+1}|\} \right\},$$

the solution u^h is explicit, and independent of h on $I_1 \cup \dots \cup I_N$:

$$u^h(t, x) = u_0(x) + t \sum_{k=0}^{N+1} \beta_{k, \ell h} \chi_k(x) \quad \text{on } [0, t_1] \times I,$$

with

$$\beta_{k, \ell h} := \begin{cases} 0, & \text{if } \alpha_{k-1} < \alpha_k < \alpha_{k+1} \text{ or if } \alpha_{k+1} < \alpha_k < \alpha_{k-1} \\ -\frac{2}{|I_k|}, & \text{if } \alpha_k > \max \{ \alpha_{k-1}, \alpha_{k+1} \} \\ \frac{2}{|I_k|}, & \text{if } \alpha_k < \min \{ \alpha_{k-1}, \alpha_{k+1} \} \end{cases} \quad (2.22)$$

for $k = 1, \dots, N$, and

$$\beta_{0, \ell h} \begin{cases} \geq 0, & \text{if } \alpha_0 < \alpha_1 \\ \leq 0, & \text{if } \alpha_0 > \alpha_1 \end{cases}$$

$$\beta_{N+1, \ell h} \begin{cases} \geq 0, & \text{if } \alpha_N > \alpha_{N+1} \\ \leq 0, & \text{if } \alpha_N < \alpha_{N+1}. \end{cases}$$

By letting $h \rightarrow 0$, we find that $u(x, t)$ remains a step function on I , its evolution is explicit on $(I_1 \cup \dots \cup I_N)$, and on I_0 and I_{N+1} it is monotonically increasing/decreasing, depending on the value on I_1 and I_N .

This formula will then continue to hold until a maximum/minimum disappear: suppose for instance that $\alpha_2 > \max\{\alpha_1, \alpha_3\}$. Then, after a certain time t'_1 , the value of $u(t)$ on I_2 becomes equal to $\max\{\alpha_1, \alpha_3\}$. Then, we simply take $u(t'_1)$ as initial data and we repeat the construction.

After repeating this at most N times, all the maxima and minima inside I disappear, and $u(t)$ is monotonically decreasing/increasing on I .

For instance, if $I = \mathbb{R}$ and the initial data is a compactly supported step function, then $u \equiv 0$ after some finite time T (which we call *extinction time*). On the other hand, if u_0 is an increasing (resp. decreasing) step function, then it will remain constant in time.

Remark 2.7 The analysis done up to now can be extended to the case of suitable initial value problems on intervals with boundary condition. For instance, the dynamic of the Dirichlet problem is analogous to the one described above for the Cauchy problem with compactly supported initial data; we leave the details to the interested reader. Let us consider next the Neumann problem on some closed interval $[a, b] = I_0 \cup \dots \cup I_{N+1}$. The dynamics on $I_1 \cup \dots \cup I_N$ is known by our analysis (which, as we observed before, is “local”). To understand the dynamics on I_0 and I_N , we go back to the time discretized problem: the Euler-Lagrange equations in this case are still (2.6), but with the additional Neumann condition $z_h(b) = z_h(a) = 0$. It is

easy to check that this last condition allows to uniquely characterize the value of u_h inside I_0 and I_{N+1} .

For example, if $u_0 = \sum_{k=1}^{N+1} \alpha_k \chi_{I_k}$ with $\alpha_1 \leq \dots \leq \alpha_{N+1}$ (i.e. u_0 is monotonically increasing), then

$$u(t) = u_0 + t \left(\frac{1}{|I_0|} \chi_{I_0} - \frac{1}{|I_{N+1}|} \chi_{I_{N+1}} \right)$$

(i.e. the value on I_0 increases, while the one on I_{N+1} decreases). This holds true until a jump disappears, and then one simply repeat the construction. We leave the details for the general case to the interested reader.

2.6 Some properties of solutions to the TVF

In this section we prove some local “regularity” properties enjoyed by solutions of the TVF (see also [13, 15]). The key fact behind these results is that the TVF is contractive in any L^q space with $q \in [1, \infty]$, cf. for example [5]. This contractivity property is not so surprising, since it holds also for the p -Laplacian for any $p > 1$. We are going to show that most of the properties which holds in the case of step functions, can be easily extended to the general case. An example is the following:

Theorem 2.8 (Local continuity) *Assume that u_0 is continuous on some open interval I . Then also the corresponding solution $u(t)$ is continuous on the same interval I and the oscillation is contractive, namely*

$$\sup_I u(t) - \inf_I u(t) =: \text{osc}_I(u(t)) \leq \text{osc}_I(u_0).$$

Proof. By the contractivity of the TVF in L^∞ , we have

$$\|u(t) - v(t)\|_\infty \leq \|u_0 - v_0\|_\infty \quad (2.23)$$

for any two given solutions $u(t), v(t)$ corresponding to initial data u_0, v_0 , with $u_0 - v_0 \in L^\infty$.

Since u_0 is continuous in I , we can find a family of functions u_0^ε such that $u_0^\varepsilon = u_0$ outside I , u_0^ε are step functions inside I , and $\|u_0 - u_0^\varepsilon\|_\infty \leq \varepsilon$. Then by (2.23) we get

$$\|u(t) - u^\varepsilon(t)\|_\infty \leq \|u_0 - u_0^\varepsilon\|_\infty \leq \varepsilon \quad \text{for all } t > 0$$

where $u^\varepsilon(t)$ is the solution to the TVF corresponding to u_0^ε , which is still a step function inside I . We now observe that the elementary inequality

$$\left| \text{osc}_I(u(t)) - \text{osc}_I(u^\varepsilon(t)) \right| \leq 2 \|u(t) - u^\varepsilon(t)\|_{L^\infty(I)} \leq 2\varepsilon \quad (2.24)$$

holds. Moreover, by looking at the explicit formulas for the evolution of $u^\varepsilon(t)$ inside I , cf. Section 2.5, it is immediate to check that $\text{osc}_I(u^\varepsilon(t))$ is decreasing in time. Hence

$$\text{osc}_I(u(t)) \leq \text{osc}_I(u^\varepsilon(t)) + 2\varepsilon \leq \text{osc}_I(u_0^\varepsilon) + 2\varepsilon \leq \text{osc}_I(u_0) + 4\varepsilon.$$

We conclude letting $\varepsilon \rightarrow 0$. \square

Remark 2.9 The above theorem still holds if u_0 is not continuous on I : in that case one has to replace sup and inf by esssup and essinf, and to prove the result one can use the comparison principle: if $u^+(t)$ and $u^-(t)$ are the solution starting respectively from

$$u^+(x) := \begin{cases} u_0(x) & \text{if } x \notin I; \\ \operatorname{esssup}_I u_0 & \text{if } x \in I; \end{cases} \quad u^-(x) := \begin{cases} u_0(x) & \text{if } x \notin I; \\ \operatorname{essinf}_I u_0 & \text{if } x \in I; \end{cases}$$

then $u^-(t) \leq u(t) \leq u^+(t)$, $u^+(t)$ and $u^-(t)$ are both constant on I , and $\|u^-(t, x) - u^-(t, x)\|_{L^\infty(I)}$ is decreasing in time. However, since we will never use this fact, we leave the details to the interested reader.

2.7 Further properties

Arguing by approximation as done in Theorem 2.8 above (using either the stability in L^∞ or simply the stability in L^1 , depending on the situation), we can easily deduce other *local properties of the TVF, valid on any subinterval I* where the solution $u(t)$ is considered (we leave the details of the proof to the interested reader):

- (i) The set of discontinuity points of $u(t)$ is contained in the set of discontinuity points of u_0 , i.e. “the TVF does not create new discontinuities”.
- (ii) The number of maxima and minima decreases in time.
- (iii) If u_0 is monotone on an interval I , then $u(t)$ has the same monotonicity as u_0 on I . Moreover, if u_0 is monotone on the whole \mathbb{R} , then it is a stationary solution to the Cauchy problem for the TVF.
- (iv) As a direct consequence of Theorem 2.8, $C^{0,\alpha}$ -regularity is preserved along the flow for any $\alpha \in (0, 1]$. (Similar results have been obtained for the denoising problem and for the Neumann problem for the TVF in [13].) Moreover, if $u_0 \in W^{1,1}(\mathbb{R})$, then $u(t) \in W^{1,1}(\mathbb{R})$ (this is a consequence of the fact that the oscillation does not increase on any subinterval).
- (v) If $u_0 \in BV_{loc}(\mathbb{R})$, a priori we do not have a well-defined semigroup. However, in this case u_0 is locally bounded and the set of its discontinuity points is countable (see [2, Section 3.2]), and so in particular has Lebesgue measure zero. Hence, by classical theorems on the Riemann integrability of functions, we can find two sequences of step functions such that $u_0^{\varepsilon,-} \leq u_0 \leq u_0^{\varepsilon,+}$ and $\|u_0^{\varepsilon,+} - u_0^{\varepsilon,-}\|_1 \leq \varepsilon$ (the number of steps will in general be infinite, but finite on any bounded interval). Then, by approximation we can still define a dynamics, which will still be contractive in any L^p space.

Behaviour near maxima and minima. We conclude this section by giving an informal description of the evolution of a general solution (excluding “pathological” cases).

Assume that u_0 has a local maximum at x_0 . Then, at least for short time, the solution is explicitly given near x_0 by

$$u(t, x) = \min\{u_0(x), h(t)\},$$

where the constant value $h(t)$ is implicitly defined by

$$\int_{I_0} [u_0(x) - h(t)]_+ dx = 2t,$$

I_0 being the connected component of $\{u_0 > h(t)\}$ containing x_0 , see Figure 4. For a minimum point the argument is analogous. The dynamics goes on in this way until a local minimum

“merges” with a local maximum, and then one can simply start again the above description starting from the new configuration.




Figure 3: *Dynamic of TVF at maximum and minimum points.*

2.8 Rescaled flow and stationary solutions

Let $u_0 \in BV(\mathbb{R}) \cap L^1(\mathbb{R})$ be a nonnegative compactly supported initial datum. First we show that u_0 extinguishes in finite time, and we calculate the explicit extinction time. (Note that, even in general dimension, estimates from above and from below on the extinction time were already known, see for example [4, 5, 18].)

Proposition 2.10 (Loss of mass and extinction time) *Let $u(t)$ be the solution to the Cauchy problem in \mathbb{R} for the TVF, starting from a non-negative compactly supported initial datum $u_0 \in L^1(\mathbb{R})$. Then the following estimates hold:*

$$\int_{\mathbb{R}} u(t, x) dx = \int_{\mathbb{R}} u_0(x) dx - 2t = 2(T - t) \quad \text{for all } t \geq 0, \quad (2.25)$$

and the extinction time for u is given by

$$T = T(u_0) = \frac{1}{2} \int_{\mathbb{R}} u_0(x) dx. \quad (2.26)$$

Proof. Arguing by approximation and using the stability of the TVF in L^1 , it suffices to consider the case when u_0 is a nonnegative step function. Assume that $\text{supp}(u_0) \subset [a, b]$ and that u_0 jumps both at a and at b . Then, by the explicit formula for $u(t)$, we immediately deduce that $u(t)$ jumps both at a and b , with $z(t, a) = 1$ and $z(t, b) = -1$ (since $u(t)$ is nonnegative as well). Hence

$$\frac{d}{dt} \int_{\mathbb{R}} u(t, x) dx = \int_a^b \partial_x z(x, t) dx = z(b) - z(a) = -2,$$

from which (2.25)-(2.26) follow. \square

Remark. Let us point out that there is no general explicit formula for the extinction time when u_0 changes sign.

The rescaled flow. We now are interested in describing the behavior of the solution near the extinction time. To this end we need to perform a logarithmic time rescaling, which maps the interval $[0, T)$ into $[0, +\infty)$, where T is the extinction time corresponding to the initial datum u_0 . We define

$$w(s, x) = \frac{T}{T-t} u(t, x), \quad Z(s, x) = z(t, x), \quad s = T \log \left(\frac{T}{T-t} \right), \quad t = T \left(1 - e^{-s/T} \right), \quad (2.27)$$

where $u(t)$ is a solution to the TVF. Then

$$\partial_s w(s, x) = \partial_x Z + \frac{w}{T}, \quad Z \cdot D_x w = |D_x w|, \quad w(0, x) = u_0(x). \quad (2.28)$$

We observe that stationary solutions $S(x)$ for the rescaled equation for w correspond to separation of variable solutions in the original variable, namely

$$-\partial_x Z = \frac{S}{T} \quad \text{provides the separate variable solution} \quad U_T(t, x) := \frac{T-t}{T} S(x).$$

We need now to characterize the stationary solutions. To this aim, we first have to define the “extended support” of a function f as the smallest interval that includes the support of f :

$$\text{supp}^*(f) = \inf \{ [a, b] \mid \text{supp}(f) \subseteq [a, b] \}.$$

Theorem 2.11 (Stationary solutions) *All compactly supported solutions of the equation*

$$-\partial_x Z = \frac{S}{T}, \quad Z \cdot D_x S = |D_x S|, \quad (2.29)$$

are of the form

$$S(x) = \frac{2T}{b-a} \chi_{[a,b]}(x), \quad (2.30)$$

with $[a, b] \subseteq \mathbb{R}$.

Proof. Let us assume that $\text{supp}^* S = [a, b]$. Since S is nonnegative we have $Z(a) = 1$, $Z(b) = -1$. We claim that $-1 < Z < 1$ on (a, b) .

Indeed, assume by contradiction that $Z(x_0) = 1$ for some point $x_0 \in (a, b)$ (the case resp. $Z(x_0) = -1$ is completely analogous). Then, using again that S is nonnegative we obtain $\partial_x Z = -\frac{S}{T} \leq 0$, which implies that $Z \equiv 1$ on $[a, x_0]$. Hence $S = -T \partial_x Z = 0$ on $[a, x_0]$, which contradicts the definition of $\text{supp}^*(S)$.

Thanks to the claim, since $Z \cdot D_x S = |D_x S|$ we easily deduce that $|D_x S|(a, b) = 0$, that is, S is constant inside (a, b) . To find the value of such a constant, we simply integrate the equation over $[a, b]$, and we get

$$\int_a^b \frac{S}{T} dx = - \int_a^b \partial_x Z(x) dx = Z(a) - Z(b) = 2.$$

and (2.30) follows. \square

Corollary 2.12 (Separate variable solutions) *All compactly supported solutions of the TVF obtained by separation of variables are of the form*

$$U_T(t, x) = 2 \frac{T-t}{b-a} \chi_{(a,b)}(x). \quad (2.31)$$

where $T > 0$ and $[a, b] \subseteq \mathbb{R}$.

Proposition 2.13 (Mass conservation for rescaled solutions) *Let $u(t)$ be the solution to the TVF corresponding to a nonnegative initial datum $u_0 \in BV(\mathbb{R}) \cap L^1(\mathbb{R})$. Let $w(s)$ be the corresponding rescaled solution, as in (2.27), then we have that*

$$\int_{\mathbb{R}} w(s, x) dx = \int_{\mathbb{R}} u_0(x) dx. \quad (2.32)$$

Proof. From (2.25), (2.27), and the fact that the extinction time is given by $2T = \int_{\mathbb{R}} u_0 dx$, we deduce that

$$\int_{\mathbb{R}} w(s, x) dx = \frac{T}{T-t} \left[\int_{\mathbb{R}} u_0(x) dx - 2t \right] = e^{s/T} \left[\int_{\mathbb{R}} u_0(x) dx - 2T \left(1 - e^{-s/T} \right) \right] = \int_{\mathbb{R}} u_0(x) dx.$$

□

Proposition 2.14 (Stationary solutions are asymptotic profiles) *Let $w(s, x)$ be a solution to the rescaled TVF corresponding to a non-negative initial datum $u_0 \in BV(\mathbb{R}) \cap L^1(\mathbb{R})$. Then there exists a subsequence $s_n \rightarrow \infty$ such that $w(s_n, \cdot) \rightarrow S$ in $L^1(I)$ as $n \rightarrow \infty$ where S is a stationary solution as in (2.30). Equivalently we have that there exists a sequence of times $t_n \rightarrow T$ as $n \rightarrow \infty$ such that*

$$\left\| \frac{u(t_n, \cdot)}{T - t_n} - \frac{S}{T} \right\|_{L^1} \xrightarrow{n \rightarrow \infty} 0.$$

where S is a stationary solution.

Proof. This is a well known result, see e.g. Theorem 4.3 of [3] or Theorem 3 of [4] for the homogeneous Dirichlet problem on bounded domains. See also the book [5]. □

Remark. From the above result we cannot directly deduce the correct extinction profile for the TVF, since there is not uniqueness of the stationary state, as Theorem 2.11 shows. Indeed a priori there may exist different subsequences such that the solution w approaches two different stationary states along the two subsequences. We shall prove in the next section that such phenomenon does not occur.

2.9 Asymptotics of the TVF

Here we want to characterize the asymptotic (or extinction) profile for solutions to the TVF in function of the non-negative initial datum $u_0 \in BV(\mathbb{R}) \cap L^1(\mathbb{R})$.

Theorem 2.15 (Extinction profile for solutions to the TVF) *Let $u(t, x)$ be a solution to the TVF corresponding to a non negative initial datum $u_0 \in BV(\mathbb{R})$ with $\text{supp}^*(u_0) = [a, b]$, and set*

$$T = \frac{1}{2} \int_a^b u_0(x) dx.$$

Then $\text{supp}(u(t)) = [a, b]$ for all $t \in (0, T)$ and

$$\left\| \frac{u(t, \cdot)}{T-t} - 2 \frac{\chi_{[a,b]}}{b-a} \right\|_{L^1([a,b])} \xrightarrow{t \rightarrow T} 0. \quad (2.33)$$

Remarks. (i) The above theorem shows two important facts: firstly, the support of the solution becomes instantaneously the “extended support” of the initial datum, which is the support of the extinction profile. Secondly, on $[a, b] = \text{supp}^*(u_0)$ we consider the quotient $u(t, x)/U_T(t, x)$, where U_T is the separate variable solution $U_T(t, x) = (T-t)S(x)$, and $S(x) = 2 \frac{\chi_{[a,b]}}{b-a}$ (see (2.30) and (2.31)). It is interesting to point out that U_T is explicitly characterized in function of the extinction time (i.e. $\frac{1}{2} \int u_0$) and of extended support of the initial datum. Then (2.33) can be rewritten as

$$\left\| \frac{u(t, \cdot)}{U_T(t, \cdot)} - 1 \right\|_{L^1([a,b])} \xrightarrow{t \rightarrow T} 0.$$

(In literature this result is usually called *convergence in relative error*.) Equivalently, L^1 -norm of the difference decays at least with the rate

$$\|u(t, \cdot) - U_T(t, \cdot)\|_{L^1(\mathbb{R})} \leq o(T-t).$$

In the next paragraph we will prove that the $o(1)$ appearing in the above rate cannot be quantified/improved, so that the convergence result of Theorem 2.15 is sharp.

(ii) The result of the above theorem can be restated in terms of the rescaled flow of Subsection 2.8: if $w(s, x)$ is the rescaled solution corresponding to $w(0, x) = u_0$ (see 2.27), then the relative error $w(s, x)/S \rightarrow 1$ as $s \rightarrow \infty$ in $L^1(\text{supp}^*(u_0))$.

Proof. The proof is divided into several steps.

• **STEP 1. Compactness estimates.** Let $z(x, t)$ be associated to the $u(t, x)$ (as in the definition of strong solution, see Section 2.3). We claim that

$$\|\partial_x z(x, t)\|_2 \leq \frac{2\|u_0\|_2}{t}. \quad (2.34)$$

The proof of this fact is quite standard in semigroup theory, once one observes that $\partial_t u(t, x) = \partial_x z_u(x, t)$. Indeed, the homogeneity of the semigroup implies that

$$\|\partial_t u(t, x)\|_p \leq \frac{2\|u_0\|_p}{t} \quad \forall p \in [1, \infty]. \quad (2.35)$$

Although the latter estimate is classical and due to Benilan and Crandall [12], we briefly recall here the proof for convenience of the reader. Since $u(t, x)$ is a solution to the TVF starting from $u(0, \cdot) = u_0(\cdot)$, it is then clear that $u_\lambda(t, x) = \lambda u(\lambda^{-1}t, x)$ is again a solution to the TVF starting from $u_\lambda(0, \cdot) = \lambda u_0(\cdot)$, for all $\lambda \geq 0$. Then

$$\begin{aligned} u(t+h, x) - u(t, x) &= \frac{t+h}{t} u_\lambda(t, x) - u(t, x) = \lambda^{-1} u_\lambda(t, x) - u(t, x) \\ &= (\lambda^{-1} - 1) u_\lambda(t, x) + (u_\lambda(t, x) - u(t, x)) \end{aligned}$$

where we have defined $\lambda := t/(t+h) > 0$. We now use the contraction property of the TVF in any L^p -space to conclude that

$$\begin{aligned} \|u(t+h, x) - u(t, x)\|_p &\leq (\lambda^{-1} - 1)\|u_\lambda(t, x)\|_p + \|u_\lambda(t, x) - u(t, x)\|_p \\ &\leq |\lambda^{-1} - 1|\|u_0\|_p + \|u_\lambda(0, x) - u(0, x)\|_p \\ &= |\lambda^{-1} - 1|\|u_0\|_p + |\lambda - 1|\|u_0\|_p = \frac{h}{t} + \frac{h}{t+h}\|u_0\|_p \leq \frac{2h}{t}\|u_0\|_p. \end{aligned}$$

Letting $h \rightarrow 0^+$, (2.35) follows.

• **STEP 2. Stability up to T^- .** Since $u_0 \in BV(\mathbb{R})$, as in Subsection 2.7, Property (v), we can find two sequences of step functions $u_0^{\varepsilon,-} \leq u_0 \leq u_0^{\varepsilon,+}$ such that $\|u_0^{\varepsilon,+} - u_0^{\varepsilon,-}\|_1 \leq \varepsilon$. In particular, by the formula for the extinction time, we deduce that

$$T - \varepsilon/2 \leq T(u_0^{\varepsilon,-}) \leq T(u_0^{\varepsilon,+}) \leq T + \varepsilon/2.$$

Moreover, up to replacing $u_0^{\varepsilon,+}$ with $\min\{u_0^{\varepsilon,+}, \|u_0\|_\infty \chi_{\text{supp}^*(u_0)}\}$, we have that

$$\text{supp}^*(u_0^{\varepsilon,-}), \text{supp}^*(u_0^{\varepsilon,+}) \rightarrow \text{supp}^*(u_0) \quad \text{as } \varepsilon \rightarrow 0.$$

Since the evolution of step function is explicit, it is immediately checked that

$$\text{supp}(u^{\varepsilon,-}(t)) = \text{supp}^*(u_0^{\varepsilon,-}), \quad \text{supp}(u^{\varepsilon,+}(t)) = \text{supp}^*(u_0^{\varepsilon,+})$$

for $t \in (0, T - \varepsilon/2]$ (indeed, for any subinterval $I \subset \subset \text{supp}^*(u_0^{\varepsilon,-})$ where $u_0^{\varepsilon,-}$ vanishes, $u^{\varepsilon,-}(t)$ becomes instantaneously positive, see Figure 3. By the parabolic maximum principle, this implies

$$\text{supp}^*(u_0^{\varepsilon,-}) \subset \text{supp}(u(t)) \subset \text{supp}^*(u_0^{\varepsilon,+}) \quad \text{for } t \in (0, T - \varepsilon/2],$$

which by the arbitrariness of ε implies that

$$\text{supp}(u(t)) = [a, b] \quad \forall t \in (0, T). \quad (2.36)$$

• **STEP 3. Convergence of z and shape of $u(t)$ before the extinction time.** By (2.36) and the fact that $u(t)$ is nonnegative, we deduce that $z(t, a) = 1$ and $z(t, b) = -1$ for all $t \in (0, T)$. Estimates (2.34) imply that $z(t, \cdot)$ is uniformly bounded in $C^{1/2}([a, b])$, hence it is compact in $C^0([a, b])$. Moreover, by (2.28) and Proposition 2.14 there exists a sequence $s_k = T \log\left(\frac{T}{T - \varepsilon_k}\right) \rightarrow \infty$ (i.e. $\varepsilon_k \rightarrow 0^+$) such that $\partial_s w(s_k, x) \rightarrow 0$. Hence, up to extracting a subsequence, we deduce that

$$\lim_{k \rightarrow \infty} z(T - \varepsilon_k) = 1 - \int_a^x \frac{S(y)}{T} dy \quad \text{uniformly on } [a, b],$$

where $S = \frac{2T}{\beta - \alpha} \chi_{[\alpha, \beta]}(x)$ is a stationary solution, with $a \leq \alpha \leq \beta \leq b$ (observe that, by Step 2, the support can only shrink).

• **STEP 4. Shape of $u(t)$ before the extinction time.** By Step 3 we know that $z(T - \varepsilon_k)$ converges uniformly on $[a, b]$ to the function

$$z_S(x) = \begin{cases} 1 & \text{if } a \leq x \leq \alpha \\ \frac{\alpha + \beta}{\beta - \alpha} - \frac{2}{\beta - \alpha}x & \text{if } \alpha < x < \beta \\ -1 & \text{if } \beta \leq x \leq b. \end{cases}$$

Hence, for k sufficiently large, there exist $\alpha < \alpha_{\varepsilon_k} < \beta_{\varepsilon_k} < \beta$ such that $-1 < z(T - \varepsilon_k) < 1$ on $[\alpha_{\varepsilon_k}, \beta_{\varepsilon_k}]$, $z(T - \varepsilon_k) > -1$ on $[a, \alpha_{\varepsilon_k}]$, $z(T - \varepsilon_k) < 1$ on $[\beta_{\varepsilon_k}, b]$, and $|\alpha - \alpha_{\varepsilon_k}| + |\beta - \beta_{\varepsilon_k}| \rightarrow 0$ as $\varepsilon \rightarrow 0$. Since $z(T - \varepsilon_k) \cdot D_x u(T - \varepsilon_k) = |D_x u(T - \varepsilon_k)|$, we easily deduce that $u(T - \varepsilon)$ is increasing on $[a, \alpha_{\varepsilon_k}]$, constant on $[\alpha_{\varepsilon_k}, \beta_{\varepsilon_k}]$ and decreasing on $[\beta_{\varepsilon_k}, b]$.

• **STEP 5. Solutions with only one maximum point.** Fix k large enough, and consider the evolution of $u(t)$ on the time interval $[T - \varepsilon_k, T]$. By Step 4 and the discussion at the end of Subsection 2.7. the evolution of $u(t)$ is explicit:

$$u(t, x) = \min\{u(T - \varepsilon_k, x), h(t)\} \quad \forall t \in [T - \varepsilon_k, T]$$

where $h(t) > 0$ is implicitly defined by

$$\int_{\mathbb{R}} [u(T - \varepsilon_k, x) - h(t)]_+ dx = 2t. \quad (2.37)$$

Since $\text{supp}(u(t)) = [a, b]$ (see Step 2), the above formula shows that the set $[a(t), b(t)]$ where $u(t)$ is constant expands in time and converges to $[a, b]$. Moreover, from (2.37) we easily obtain the estimate

$$-\frac{2}{b-a} \leq \dot{h}(t) \leq -\frac{1}{b-a}$$

for t close to T . Hence, the equation $\partial_t u = \partial_x z$ implies that

$$-\frac{2}{b-a} \leq \partial_x z(t) \leq -\frac{1}{b-a} \quad \text{on } [a(t), b(t)]. \quad (2.38)$$

Since $[a(t), b(t)] \rightarrow [a, b]$, the uniform convergence of $z(t)$ to z_S is compatible with (2.38) if and only if

$$z_S(x) = \frac{a+b}{b-a} - \frac{2}{b-a}x \quad \text{on } [a, b],$$

i.e. the unique possible limiting profile is $2T \frac{\chi_{[a,b]}}{b-a}$, as desired. \square

2.10 Rates of convergence

After proving asymptotic convergence to a stationary state, the next natural question is whether there exists a universal rate of convergence to it. As the next theorem shows, the answer is negative.

Before stating the result, let us make precise what do we mean by decay rate.

Definition 2.16 Let $\xi : [0, \infty) \rightarrow [0, \infty)$ be a continuous increasing function, with $\xi(0) = 0$. We say that ξ is a rate function if, for any solution $u(t)$ of the TVF,

$$\left\| \frac{u(t)}{T-t} - \frac{S}{T} \right\|_{L^1(I)} \leq \xi(T-t) \quad \text{for any } t \text{ close to the extinction time } T. \quad (2.39)$$

The following result shows that there cannot be a universal rate of convergence to any stationary profile.

Theorem 2.17 (Absence of universal convergence rates) *For any rate function $\xi : [0, \infty) \rightarrow [0, \infty)$, there exists an initial datum $u_0 \in BV(\mathbb{R})$, with $\text{supp}^*(u_0) = [0, 1]$, such that*

$$2\xi(T-t) \leq \left\| \frac{u(t)}{T-t} - 2\chi_{[0,1]} \right\|_{L^1(I)}, \quad \text{for any } 0 \leq T-t \leq 1. \quad (2.40)$$

Proof. Let us fix a rate function $\xi : \mathbb{R} \rightarrow \mathbb{R}$. It is not restrictive to assume that ξ is strictly increasing, and that $\xi(s) \geq s$ for any $s \in [0, 1]$.

Let $\xi^{-1} = [0, \infty) \rightarrow [0, \infty)$ be the inverse of ξ , so that $\xi^{-1}(s) \leq s$ for any $s \in [0, 1]$, and choose the initial datum u_0 to be

$$u_0(x) = \begin{cases} c_0 \xi^{-1}(x), & \text{if } 0 \leq x \leq \frac{1}{4} \\ 1, & \text{if } \frac{1}{4} < x < \frac{3}{4} \\ c_0 \xi^{-1}(1-x), & \text{if } \frac{3}{4} \leq x \leq 1 \end{cases} \quad (2.41)$$

with $c_0 := 1/\xi^{-1}(1/4)$ (so that $u_0(1/4) = u_0(3/4) = 1$). By Theorem 2.15 we know that the solution $u(t)$ corresponding to u_0 extinguish at time

$$\frac{1}{4} \leq T = \frac{1}{2} \int_0^1 u_0(x) \, dx \leq \frac{1}{2}$$

and that $u(t)/(T-t)$ converges strongly in $L^1([0, 1])$ to $S/T = 2\chi_{[0,1]}$ as $t \rightarrow T$. First we prove that the L^∞ -norm satisfies the bound

$$2(T-t) \leq \|u(t)\|_\infty \leq 4(T-t). \quad (2.42)$$

The first equality follows by the loss of mass formula (2.25) and Hölder inequality on $[0, 1]$:

$$2(T-t) = \|u(t)\|_1 \leq \|u(t)\|_\infty.$$

The second inequality follows as we know the explicit behaviour of the solution around maximum points (see the end of Subsection 2.7), namely

$$u(t, x) = \min\{u_0(x), h(t)\}$$

where $h(t) > 0$ satisfies

$$\int_0^1 [u_0(x) - h(t)]_+ \, dx = 2t. \quad (2.43)$$

Then $u(t)$ is constant on an interval of the form $[\alpha(t), \beta(t)]$, with $\alpha(t) \rightarrow 0^+$ and $\beta(t) \rightarrow 1^-$, see Figure 4.

Hence

$$2(T-t) = \|u(t)\|_1 = \int_0^1 u(t, x) \, dx \geq \int_{1/2}^{3/4} u(t, x) \, dx = \frac{1}{2} \|u(t)\|_\infty,$$

and so $\|u(t)\|_\infty \leq 4(T-t)$, and inequality (2.42) is proved.

Now, let $\alpha_0(t) \in [0, \alpha(t)]$ and $\beta_0(t) \in [\beta(t), 1]$ be the unique points such that $u(t, \alpha_0(t))/(T-t) = u(t, \beta_0(t))/(T-t) = 2$ (such points exists thanks to the lower bound $\|u(t)\|_\infty \geq 2(T-t)$, see also Figure 4). Let \mathbb{T}_{α_0} (resp. \mathbb{T}_{β_0}) be the rectangular triangle with height $\{0\} \times [0, 2]$




Figure 4: **Left:** *Dynamic of $u(t)$:* black: $u_0(x)$, blue: $u(t, x)$, red: $u(t + h)$. **Right:** *Rescaled dynamic:* black: $u_0(x)$ (dashdot) and u_0/T (cont.), blue: $S(x) = 2\chi_{[0,1]}$, red: $u(t, x)/(T - t)$, green: definition of $\alpha_0(t), \beta_0(t)$ and $\mathbb{T}_{\alpha_0}, \mathbb{T}_{\beta_0}$.

(resp. $\{1\} \times [0, 2)$) and basis $[0, \alpha(t)] \times \{2\}$ (resp. $[\beta(t), 1] \times \{2\}$), as depicted in Figure 4. Denote by $|\mathbb{T}_{\alpha_0}|, |\mathbb{T}_{\beta_0}|$ their measure. Then, since $\xi^{-1}(s) \geq s$ we easily obtain the estimate

$$\begin{aligned}
& \left\| \frac{u(t)}{T-t} - \frac{S}{T} \right\|_{L^1([0,1])} = \int_0^1 \left| \frac{u(t, x)}{T-t} - 2\chi_{[0,1]}(x) \right| dx \\
&= \int_0^{\alpha(t)} \left| \frac{u(t, x)}{T-t} - 2 \right| dx + \int_{\alpha(t)}^{\beta(t)} \left| \frac{u(t, x)}{T-t} - 2 \right| dx + \int_{\beta(t)}^1 \left| \frac{u(t, x)}{T-t} - 2 \right| dx \\
&\geq \int_0^{\alpha(t)} \left| \frac{u(t, x)}{T-t} - 2 \right| dx + \int_{\beta(t)}^1 \left| \frac{u(t, x)}{T-t} - 2 \right| dx \\
&\geq \int_0^{\alpha_0(t)} \left(2 - \frac{u(t, x)}{T-t} \right) dx + \int_{\beta_0(t)}^1 \left(2 - \frac{u(t, x)}{T-t} \right) dx \\
&\geq |\mathbb{T}_{\alpha_0}| + |\mathbb{T}_{\beta_0}| = 2\alpha_0(t).
\end{aligned}$$

To estimate $\alpha_0(t)$ from below, we observe that on $[0, \alpha(t)]$ we have that $u(t, \alpha_0(t)) = u_0(\alpha_0(t)) = c_0 \xi^{-1}(\alpha_0(t))$, so

$$\alpha_0(t) = \xi \left(\frac{2}{c_0} (T-t) \right).$$

Now, recalling that ξ is strictly increasing and $\xi(1) \geq 1$, we get $\xi(2) \geq 1/4$, or equivalently $2/c_0 = 2/\xi^{-1}(1/4) \geq 1$. Hence $\alpha_0(t) \geq \xi(T-t)$, which concludes the proof. \square

Remark. The above Theorem shows that there cannot be universal rates of convergence. A similar construction will provide (nontrivial) initial data for which the convergence is as fast as desired.

Theorem 2.18 (Fast decaying initial data) *For any rate function $\xi : [0, \infty) \rightarrow [0, \infty)$, there exists an initial datum $u_0 \in L^1(I)$ such that the corresponding solution $u(t)$ satisfies*

$$\left\| \frac{u(t)}{T-t} - 2\chi_{[0,1]} \right\|_{L^1(I)} \leq \xi(8(T-t)), \quad \text{for any } 0 \leq T-t \leq 1. \quad (2.44)$$

Proof. Fix a rate function $\xi : \mathbb{R} \rightarrow \mathbb{R}$, which is continuous, increasing, $\xi(0) = 0$, and $\xi(s) \leq s$.

Let $\xi^{-1} = [0, \infty) \rightarrow [0, \infty)$ denote its inverse, and define u_0 as in (2.41), see Figure 5. Then an analysis analogous to the one done in the previous Theorem proves the result. We leave the details to the interested reader. \square




Figure 5: **Left:** *Dynamic of $u(t)$:* black: $u_0(x)$, blue: $u(t, x)$, red: $u(t+h, x)$. **Right:** *Rescaled dynamic:* black: $u_0(x)$, blue: $S(x) = 2\chi_{[0,1]}$, red: $u(t, x)/(T-t)$, green: definition of $\alpha_0(t)$, $\beta_0(t)$ and \mathbb{T}_{α_0} , \mathbb{T}_{β_0} .

3 Solutions to the SFDE and solutions to the TVF

As explained in the introduction, TVF and SFDE are formally related by the fact that “ u solves the TVF if and only if $D_x u$ solves the SFDE”.

In order to make this rigorous, we need first to explain what do we mean by a solution of the SFDE, and then we will prove the above relation by approximating the TVF with the p -Laplacian and the SFDE by the porous medium equation.

The notion of solution we consider for the SFDE is the one of mild solution. More precisely, since the multivalued graph of the function $r \mapsto \text{sign}(r)$ is maximal monotone, by the results of Benilan and Crandall [11], there exists a continuous semigroup $S_0^t : L^1(\mathbb{R}) \rightarrow L^1(\mathbb{R})$ such that $S_t^0 v_0 \in C([0, \infty); L^1(\mathbb{R}))$ is a mild solution of the SFDE. To be more precise, let φ be a maximal monotone graph in \mathbb{R} (see [10]) and consider the problem

$$\begin{cases} u_t = \Delta \varphi(u), & \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}) \\ u(0, x) = u_0(x), & x \in \mathbb{R} \end{cases} \quad (3.1)$$

where the first equation is meant in the sense that

$$u_t = \Delta w \quad \text{in } \mathcal{D}'((0, \infty) \times \mathbb{R}), \quad \text{with } w(t, x) \in \varphi(u(t, x)) \quad \text{a.e. } t, x \in \mathbb{R}. \quad (3.2)$$

We now recall the celebrated results of Benilan and Crandall [11] adapted to our setting, namely it is sufficient to consider $\varphi(r) = \text{sign}(r)$.

Theorem 3.1 (Benilan-Crandall, [11]) *Under the running assumptions, the following results hold true:*

(i) *There exists a unique solution $u \in C([0, \infty); L^1(\mathbb{R})) \cap L^\infty([0, \infty) \times \mathbb{R})$ corresponding to the*

initial datum $u_0 \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ such that (3.1) and (3.2) are satisfied.

(ii) Let $u_n \in C([0, \infty); L^1(\mathbb{R})) \cap L^\infty([0, \infty) \times \mathbb{R})$ be solutions of (3.2) corresponding to the sequence $\varphi_n : \mathbb{R} \rightarrow \mathbb{R}$, $n = 1, 2, \dots$ of maximal monotone graphs in \mathbb{R} . Assume that $0 \in \varphi_n(0)$,

$$\lim_{n \rightarrow \infty} \varphi_n(r) = \varphi(r) \quad \text{for } r \in \mathbb{R}, \quad \text{and} \quad \lim_{n \rightarrow \infty} \|u_{0,n} - u_0\|_{L^1(\mathbb{R})} = 0. \quad (3.3)$$

Then $u_n \rightarrow u$ in $C([0, \infty); L^1(\mathbb{R}))$, where u is the solutions of (3.2) corresponding to φ .

Now, let S_t^m be the semigroup associated to the FDE equation

$$\partial_t v = \Delta(v^m).$$

Since the graphs of the function $r \mapsto r^m := |r|^{m-1}r$ converge to the graph of $r \mapsto \text{sign}(r)$, we can use Theorem 3.1 with the simple choice $\varphi_n(r) = |r|^{\frac{1}{n}-1}r$ to guarantee that we have convergence (indeed as $m \rightarrow 0^+$) of $S_t^m v_0$ to $S_t^0 v_0$ in $C([0, \infty); L^1(\mathbb{R}))$, for any initial datum $v_0 \in L^1$.

On the other hand, we can consider the p -Laplacian semigroup T_t^p for $p = 1 + m$. It is well known (see for example [10] chap. 4, or [5] chap. 5) that if $u_0 \in W^{1,p}(\mathbb{R})$, then $T_t^p u_0 \in W^{1,p}(\mathbb{R})$, so that as $p \rightarrow 1^+$, strong solutions to the p -Laplacian converge to strong solutions to the TVF. (A detailed proof of this fact in dimension $n \geq 1$ can be found for instance in [4], pg. 138-142, in the framework of the Dirichlet problems on bounded domains, but it can be easily adapted to other problems, including the Cauchy one.) Hence, if u_0 is a smooth compactly supported function, $T_t^p u_0 \rightarrow T_t^1 u_0$ in $C([0, \infty); L^1(\mathbb{R}))$ as $p \rightarrow 1^+$, where T_t^1 denotes the TVF-semigroup.

Moreover, if $p = 1 + m$, we have that $\partial_x(T_t^p u_0)$ solves (in the distributional and semigroup sense) the FDE with initial datum $\partial_x u_0$, i.e. $\partial_x(T_t^p u_0) = S_t^m(\partial_x u_0)$. Hence, by letting $m \rightarrow 0^+$, we recover such a relation in the limit $p = 1$ and $m = 0$ by what we said above. We can summarize this discussion in the following:

Theorem 3.2 *Assume u_0 is a smooth compactly supported function. Let $1 < p \leq 2$, and $m = p - 1$. Then the following diagram is commutative:*

$$\begin{array}{ccc} T_t^p u_0 \in W^{1,p}(\mathbb{R}) & \xrightarrow{p \rightarrow 1^+} & T_t^1 u_0 \in W^{1,1}(\mathbb{R}) \\ \partial_x \downarrow & & \downarrow \partial_x \\ S_t^m(\partial_x u_0) \in L^{1+m}(\mathbb{R}) & \xrightarrow{m \rightarrow 0^+} & S_t^0(\partial_x u_0) \in L^1(\mathbb{R}). \end{array}$$

Note that the convergence is meant in the sense of distributions

At this point it is worth noticing that the vector field z associated to the solution u of the TVF as in Section 2.2 and the function w associated to the solutions to the SFDE as in (3.2) are the same (just by letting $v_0 = \partial_x u_0$ and $v = \partial_x u$)

$$\partial_t u = \partial_x z \quad \xrightarrow{\partial_x} \quad \partial_t v = \partial_t \partial_x u = \partial_{xx} z = \partial_{xx} w.$$

Measures as initial data. Once the correspondence between TVF and SFDE is established for smooth initial data, by stability in L^1 of both semigroups it immediately extends to $u_0 \in$

$W^{1,1}(\mathbb{R})$, and then by approximation to $BV(\mathbb{R}) \cap L^1(\mathbb{R})$ initial data. However, at the level of the SFDE this would correspond to finite measures v_0 such that $\int_{-\infty}^x v_0(dy) \in L^1(\mathbb{R})$, which is possible if and only if $\int_{-\infty}^{+\infty} v_0(dy) = 0$. Actually, this class of data correspond exactly to the one for which there is extinction in finite time (as this is the case for L^1 initial data to the TVF).

To remove this unnatural constraint on v_0 , we observe that, by Subsection 2.7, Property (v), the TVF defines a contractive semigroup also on initial data which are only in $BV_{loc}(\mathbb{R})$. In particular, the TVF is well-defined on data of the form $u_0(x) = \int_{-\infty}^x v_0(dy)$, where v_0 is a (locally) finite measure on \mathbb{R} . Hence, this allows to define measure valued solutions of the SFDE as $\partial_x T_t^1(u_0)$, and this notion coincides with the one that one would get by considering weak* limit of L^1 solutions.

Summing up, we have shown that:

- If $v_0 \in L^1(\mathbb{R})$, the unique mild solution of the SFDE of Theorem 3.1 is given by

$$S_t^0 v_0 = \partial_x \left(T_t^1 \left(\int_{-\infty}^x v_0(dy) \right) \right). \quad (3.4)$$

- Using (3.4) we can uniquely extend the generator S_t^0 to measure initial data (actually, since the semigroup T_t^1 is well-defined on $L^2(\mathbb{R})$, one could even extend the SFDE to distributional initial data in $W^{-1,2}(\mathbb{R})$).

3.1 The 1-dimensional SFDE

By (3.4), the dynamics of the SFDE can be completely recovered from the one of the TVF. We begin by illustrating some basic properties, and then, instead of trying to give a complete description of the evolution (which, by the analysis of the TVF done in the previous sections, would just be a tedious exercise), we prefer to briefly illustrate the evolution of solutions of the SFDE in some simple but representative situations. As pointed out in the previous section, we are allowed to consider measures as initial data for the SFDE, keeping in mind that the distributional x -derivative of a solution to the TVF is a solution to the SFDE.

In the same way as step functions allowed us to understand the dynamics of the TVF, we start by considering the dynamics of sum of delta masses, which is directly deduced from the one of step functions for the TVF.

Example 1. Delta masses as initial data. Let us assume that $v_0 = \sum_{i=1}^N a_i \delta_{x_i}$, with $x_1 < \dots < x_N$. Then, for $t > 0$ small (the smallness depending on the size of $|a_i|$)

$$v(t) = \sum_{i=1}^N a_i(t) \delta_{x_i},$$

with $a_i(0) = a_i$ and

$$a_i(t) = \begin{cases} a_i, & \text{if } \text{sign}(a_{i-1}) = \text{sign}(a_{i+1}) = \text{sign}(a_i) \\ \text{sign}(a_i)(|a_i| - 4t)_+, & \text{if } \text{sign}(a_{i-1}) = \text{sign}(a_{i+1}) = -\text{sign}(a_i) \\ \text{sign}(a_i)(|a_i| - 2t)_+, & \text{if } \text{sign}(a_{i-1}) \text{sign}(a_{i+1}) = -1, \end{cases} \quad (3.5)$$




Figure 6: black: $v_0(x)$, blue: $v(t, x)$, $t < t_0$, red: $v(t_1, x)$ is a stationary state.

where we use the convention $\text{sign}(a_0) := \text{sign}(a_1)$ and $\text{sign}(a_{N+1}) := \text{sign}(a_N)$. This formula holds true until one mass disappear at some time $t'_1 > 0$, and then it suffices to $v(t'_1)$ as initial data and repeat the construction (compare with Subsection 2.5).

We observe that if all a_i have the same sign, then v_0 produces a stationary solution $v(t, x) = v_0(x)$. Moreover, the total mass is conserved under the dynamics. In particular, $v(t)$ extinguishes in finite time if and only if it has zero mean, i.e. $\sum_i a_i = 0$ (however, there is no simple formula for the extinction time, see the remark after Proposition 2.10).

General properties. Arguing by approximation (or again using the direct relation with the TVF), as a consequence we have the following properties of the SFDE flow:

(i) *Nonnegative initial data.* Let $v_0 \geq 0$ be a locally finite measure, and define $u_0(x) := \int_{-\infty}^x v_0(dy) \geq 0$. Since u_0 is monotone non-decreasing, it does not evolve under the TVF, cf. (iii) in Section 2.7. Hence v_0 is a stationary solution to the SFDE. (Actually, since monotone profiles are the only stationary state for the TVF, the only stationary solutions for the SFDE are nonnegative/nonpositive initial data.)

(ii) *Only zero mean valued initial data extinguish in finite time.* If v_0 is a finite measure, $v(t)$ converges in finite time to a stationary solution \bar{v} such that $\int_{\mathbb{R}} \bar{v}(dy) = \int_{\mathbb{R}} v_0(dy)$. Moreover, $\bar{v} \equiv 0$ (i.e. v_0 extinguish in finite time) if and only if $\int_{\mathbb{R}} v_0(dy) = 0$.

Example 2. Interaction between a delta and a continuous part.

Let $v_0 = \hat{v}_0 + \alpha\delta_0$ where $\hat{v}_0 \in C(\mathbb{R})$ is positive exception made for an interval $[0, x_0]$ as depicted in Figure 6. The zero set of \hat{v}_0 is $Z(\hat{v}_0) = [0, a] \cup \{x_0\}$, and a Dirac's delta with mass α is put at $x = 0$. We assume α is much smaller than $\int_a^{x_0} |\hat{v}_0|$. The evolution basically changes at two steps:

- $0 \leq t \leq t_0$. The delta mass starts to lose its mass α by a factor $2t$ until time $t_0 = \alpha/2$ (when it extinguish). This mass is compensated by a “gain of mass” of \hat{v}_0 : its zero set near

a starts to move to the right, and at time t it is at a position $z_1(t)$ characterized by the fact that $\int_a^{a+z_1(t)} |\hat{v}_0| = 2t$ (cf. the blue area $2t$ in the Figure 6). On the other hand, the isolated point $x_0 \in Z(v_0)$ starts to “expand”, creating a zero set $[z_2(t), z_3(t)] \in Z(u(t))$, with $\int_{z_2(t)}^{x_0} |\hat{v}_0| = \int_{x_0}^{z_3(t)} |\hat{v}_0| = 2t$. This expansion of the zero set corresponds to the creation of flat parts at the level TVF (see Figure 3).

- $t_0 \leq t \leq t_1$. At time t_0 the delta disappears, and $u(t_0, x)$ is a piecewise continuous function on \mathbb{R} which is positive outside $[0, z_3(t_0)]$, it is zero on $Z(u(t_0, x)) = [0, z_1(t_0)] \cup [z_2(t_0), z_3(t_0)]$, and is negative on $(z_1(t_0), z_2(t_0))$. Starting from this time, the free boundary expands on both components in the same way as described above, until some time $t_1 > t_0$ when $z_1(t_1) = z_2(t_1)$. Observe that the loss of mass is such that $B_0 = B_1 + B_2$ (see Figure 6).

- *Reaching the stationary state in finite time.* Since $u(t_1) \geq 0$, the solution becomes stationary and $u(t) = u(t_1)$ for $t \geq t_1$.

3.2 SFDE vs LFDE.

Let us go back to the fast diffusion equation (1.1), and assume that $v_0 \geq 0$ (so $v(t) \geq 0$ for all $t \geq 0$). We remark that, changing the time scale $t \mapsto mt$, the above equation can be written in two different ways which lead to two different limiting equations: more precisely, setting $\rho(t, x) = v(t/m, x)$,

$$\begin{array}{ccc} \partial_t v = \Delta(v^m) & \xrightarrow{m \rightarrow 0^+} & \partial_t v = \Delta(\text{sign}(v)) \\ \partial_t \rho = \text{div}(\rho^{m-1} \nabla \rho) & \xrightarrow{m \rightarrow 0^+} & \partial_t \rho = \text{div}(\rho^{-1} \nabla \rho) = \Delta(\log(\rho)). \end{array}$$

Observe that the evolution of $\rho(t, x)$ on the time interval $[0, T]$ corresponds to the evolution of $v(t, x)$ on the larger time interval $[0, T/m]$, and the diffusion of v is slower than the diffusion of ρ by a factor $1/m$. So, when analyzing the limit as $m \rightarrow 0^+$, one gets two different limits, and the evolution of $\rho(t, x)$ on the time interval $0 \leq t \leq T$ corresponds to the evolution of $v(t, x)$ on the time interval $0 \leq t < \infty$ for every $T > 0$. This means that the solution to the SFDE corresponds to an evolution “infinitely slower” than the solution to the LFDE.

We saw that the Cauchy problem for the SFDE gives rise to a trivial dynamics on non-negative initial data. However, the problem becomes non-trivial if we consider for instance the Dirichlet problem for the SFDE on a closed interval I with zero boundary conditions. Indeed, by “integrating in space” such a solution, we obtain a solution to the TVF on I with Neumann boundary conditions, whose dynamics on step functions has been described in Remark 2.7. In particular, from the example given there, we can explicitly find the dynamics of a finite sum of positive deltas: if $v_0 = \sum_{i=1}^N a_i \delta_{x_i}$ with $a_i > 0$ and $x_1 \leq \dots \leq x_N$, then $v(t) = v_0 - t[\delta_{x_1} + \delta_{x_N}]$ until one delta disappears, and then one simply restarts from there. By approximation, we see that positive initial data extinguish in finite time, and the extinction time is given by $\frac{1}{2} \int_{\mathbb{R}} v_0(dy)$.

This fact and the above discussion suggest that solutions to the LFDE should extinguish instantaneously, i.e. $\rho(t) \equiv 0$ for any $t > 0$. As one can deduce by the results of Rodriguez and Vázquez [22], this is actually the case. (See also [14, 21, 22] and the book [25] for a complete theory of the logarithmic diffusion equation in one space dimension for positive initial data.) So the above discussion is correct, and gives an heuristic explanation for this immediate extinction phenomenon.

References

- [1] L. Ambrosio, V. Caselles, S. Masnou, J.-M. Morel, *Connected components of sets of finite perimeter and applications to image processing*, J. Europ. Math. Soc., **3** (2001), 39–92.
- [2] L. Ambrosio, N. Fusco, D. Pallara, *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [3] F. Andreu, C. Ballester, V. Caselles, *A parabolic Quasilinear problem for Linear Growth Functionals*, Rev. Mat. Iberoamericana **18** (2002), 135–185.
- [4] F. Andreu, V. Caselles, J. I. Diaz, J. M. Mazon, *Some qualitative properties for the total variation flow*, J. Funct. Anal., **188** (2002), no.2, 516–547.
- [5] F. Andreu, V. Caselles, J. M. Mazon, *Parabolic quasilinear equations minimizing linear growth functionals*, Progress in Mathematics, **223**, Birkhuser Verlag, Basel, 2004. xiv+340 pp. ISBN: 3-7643-6619-2.
- [6] G. I. Barenblatt, J. L. Vázquez, *Nonlinear Diffusion and Image Contour Enhancement*, Interf. and Free Bound., **6** (2004), 31–54.
- [7] G. Bellettini, V. Caselles, M. Novaga, *The total variation flow in \mathbb{R}^N* , J. Differential Equations **184** (2002), no. 2, 475–525.
- [8] G. Bellettini, V. Caselles, M. Novaga, *Explicit Solutions of the Eigenvalue Problem $\operatorname{div} \left(\frac{Du}{|Du|} \right) = u$ in \mathbb{R}^2* , SIAM J. Math. Anal. **36** (2005), 1095–1129.
- [9] V. Caselles, A. Chambolle, M. Novaga, *The Discontinuity Set of Solutions of the TV Denoising Problem and Some Extensions*, Multiscale Model. Simul. **6** (2007), 879–894.
- [10] H. Brezis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*. (French) North-Holland Mathematics Studies, No. 5. Notas de Matemática (50). North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, (1973). vi+183 pp.
- [11] P. Benilan, M. G. Crandall, *The continuous dependence on φ of solutions of $u_t - \Delta\varphi(u) = 0$* Indiana Univ. Math. J. **30** (1981), no. 2, 161–177.
- [12] P. Benilan, M. G. Crandall, *Regularizing effects of homogeneous evolution equations*, Contributions to Analysis and Geometry, (suppl. to Amer. J. Math.) (1981), Johns Hopkins Univ. Press, Baltimore, Md., pp. 23-39.
- [13] V. Caselles, A. Chambolle, M. Novaga *Regularity for solutions of the total variation denoising problem*, Rev. Mat. Iberoamericana **27**, (2011), no. 1, 233–252.
- [14] J. R. Esteban, A. Rodriguez, J. L. Vázquez, *A nonlinear heat equation with singular diffusivity*, Comm. Partial Differential Equations **13** (1988), no. 8, 985-1039.
- [15] M.-H. Giga, Y. Giga, *Evolving graphs by singular weighted curvature*, Arch. Rational Mech. Anal. **141** (1998), 117–198

- [16] M.-H. Giga, Y. Giga, *Very singular diffusion equations: second and fourth order problems*, Japan J. Indust. Appl. Math., **27** (2010), 323–345.
- [17] M.-H. Giga, Y. Giga, R. Kobayashi, *Very singular diffusion equations*, *Proc. of the last Taniguchi Symposium*, (eds. M. Maruyama and T. Sunada), Advanced Studies in Pure Math., **31** (2001), 93–125.
- [18] Y. Giga, R. Kohn, *Scale-invariant extinction time estimates for some singular diffusion equations*, Discr. Cont. Dyn. Sys., **30**, no. 2 (2011), 509–535.
- [19] K. Kielak, P. Boguslaw Mucha, P. Rybka, *Almost classical solutions to the total variation flow*, Preprint 2011, arXiv:1106.5369v1
- [20] R. Iagar, A. Sánchez, J. L. Vázquez, *Radial equivalence for the two basic nonlinear degenerate diffusion equations*, J. Math. Pures Appl., **89** (2008), no. 1, 1–24.
- [21] A. Rodríguez, J. L. Vázquez, *A well-posed problem in singular Fickian diffusion*. *Arch. Rational Mech. Anal.* **110** (1990), no. 2, 141-163.
- [22] A. Rodríguez, J. L. Vázquez, *Obstructions to existence in fast-diffusion equations*, J. Differential Equations **184** (2002), no. 2, 348-385.
- [23] L. I. Rudin, S. Osher, E. Fatemi, *Nonlinear total variation based noise removal algorithms*, Physica D, **60**, (1992), 259-268.
- [24] J. L. Vázquez, *Two nonlinear diffusion equation with finite speed of propagation*, Problems involving change of type (Stuttgart, 1988), 197–206 Lecture Notes in Phys., 359, Springer, Berlin, 1990.
- [25] J. L. Vázquez, *Smoothing and decay estimates for nonlinear diffusion equations*, vol. 33, Oxford Lecture Notes in Maths. and its Applications, Oxford Univ. Press, 2006.
- [26] J. L. Vázquez, *The Porous Medium Equation. Mathematical Theory*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2007.