

## Hoja de problemas 1: LAPLACE AND POISSON EQUATIONS

1. Prove that Laplace equation  $\Delta u = 0$  is invariant under rotations: let  $O$  be an orthogonal matrix  $n \times n$  and define

$$v(x) := u(Ox), \quad x \in \mathbb{R}^N.$$

Show that  $\Delta v = 0$ .

2. Let  $u$  be an harmonic function and let  $\phi : \mathbb{R} \mapsto \mathbb{R}$  be a smooth convex function. Prove that  $v := \phi(u)$  is a subharmonic function.

3. Show that  $x \mapsto \log |x|$  is a subharmonic function in the domain  $\mathbb{R}^N \setminus \{0\}$  if  $N \geq 2$ .

4. Show that  $v := |Du|^2$  a subharmonic function if  $u$  is harmonic.

5. Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain and  $u \in C^2(\Omega) \cap C(\bar{\Omega})$  a solution to

$$\Delta u = -1 \quad \text{en } \Omega, \quad u|_{\partial\Omega} = 0.$$

Prove that  $\forall x_0 \in \Omega$  we have that

$$u(x_0) \geq \frac{1}{2N} \min_{x \in \partial\Omega} |x - x_0|^2.$$

6. Let  $u$  be a classical solution to

$$-\Delta u = f \quad \text{en } B_1(0), \quad u = g \quad \text{en } \partial B_1(0).$$

Show that there exists a constant  $C > 0$ , independent of  $u$ , such that

$$\max_{B_1(0)} |u| \leq C (\max_{\partial B_1(0)} |g| + \max_{B_1(0)} |f|).$$

7. Let  $u$  be a positive harmonic function in  $B_r(0)$ . Use Poisson formula to show that

$$r^{N-2} \frac{r - |x|}{(r + |x|)^{N-1}} u(0) \leq u(x) \leq r^{N-2} \frac{r + |x|}{(r - |x|)^{N-1}} u(0).$$

This is an explicit form of the Harnack inequality.

8. Consider the problem

$$\begin{cases} \Delta u(x) + c(x)u(x) = 0, & x \in \Omega, \\ u(x) = g(x), & x \in \partial\Omega, \end{cases}$$

where we assume  $c(x) < 0$ . Show that this problem has a unique solution. Show by an example that when  $c(x) > 0$  uniqueness fails.

9. (Schwartz Reflection Principle) Consider the open semiball  $U^+ = \{x \in \mathbb{R}^N : |x| < 1, x_N > 0\}$ . Let  $u \in C^2(U^+)$  be harmonic in  $U^+$  with  $u = 0$  on  $\partial U^+ \cap \{x_N = 0\}$ . Given  $x \in U = B_1(0)$  we define

$$v(x) := \begin{cases} u(x) & \text{si } x_N \geq 0, \\ -u(x_1, \dots, x_{n-1}, -x_N) & \text{si } x_N < 0. \end{cases}$$

Show that  $v$  is harmonic in  $U$ .

10. Let  $\Omega \subset \mathbb{R}^N$ , be a domain,  $N \geq 2$ , and  $x_0 \in \Omega$ . Let  $u$  be a *bounded* harmonic function in  $\Omega_0 := \Omega \setminus \{x_0\}$ . Show that we can define a value  $u(x_0)$  such that the extended function is harmonic on the whole  $\Omega$ .

11. Let  $\Omega \subset \mathbb{R}^N$ , be a bounded domain,  $N \geq 2$ , and let  $x_0 \in \Omega$ . Define  $\Omega_0 := \Omega \setminus \{x_0\}$  and let  $u$  and  $v$  be two harmonic functions in  $\Omega_0$ , continuous in  $\Omega_1 = \Omega_0 \cup \partial\Omega$  and such that: (i)  $u(x) \leq v(x)$  for all  $x \in \partial\Omega$ ; (ii)  $|u(x)| \leq M$ ,  $|v(x)| \leq M$  for all  $x \in \Omega_1$ . Use the Maximum Principle to show that  $u(x) \leq v(x)$  for all  $x \in \Omega_1$ .
12. Find an expression for the Green function of the Dirichlet problem for the Laplace equations in an annular region  $B_R(x_0) \setminus B_r(x_0)$ , with  $0 < r < R$ .
13. Show that a solution to  $\Delta u - u^2 = 0$  in a domain  $\Omega$  cannot attain its maximum in  $\Omega$ , except if  $u \equiv 0$ .
14. Let  $u \in C^2(B_1(0)) \cap C(\overline{B_1(0)})$  be a solution to the Dirichlet problem

$$\begin{cases} \Delta u = u^2 + f(|x|), & x \in B_1(0), \\ u(x) = 1, & x \in \partial B_1(0), \end{cases}$$

where  $f(|x|) \geq 0$  is of class  $C^1(\Omega)$ . Calculate the maximum of  $u$  in  $\overline{B_1(0)}$  and show that it does not depend on  $f$ .